

FAMILY CAPSIDÆ.

387. *MIRIS PUNCTULATUS* Kirby.—Length of body 3 lines. Two specimens taken in Lat. 65°.

[278.] Body oblong, pale, somewhat lurid with a very slight tint of green. Head triangular, impunctured; eyes a little embrowned, prominent; antennæ as long as the prothorax, substestaceous with the two last joints and the underside of the first black; thorax, scutellum and hemelytra very minutely punctured; a small portion of the prothorax adjoining the head is separated from the rest by an impressed sinuous line and is not punctured; legs pale; abdomen subpyramidal, black above, pale underneath.

388. *MIRIS VENTRALIS* Kirby.—Length of body 3 lines. Taken with the preceding.

Very similar to *M. punctulatus*, and perhaps only a variety, but the hemelytra are faintly clouded with black, the underside of the abdomen is dusky with two longitudinal reddish spots or stripes.

ENTOMOLOGICAL CLUB OF THE AMERICAN ASSOCIATION
FOR THE ADVANCEMENT OF SCIENCE.

The Annual Meeting of the Club will be held at St. Louis, Mo., on Tuesday, the 20th of August, 1878, at 3 o'clock, p. m. All Entomologists who are interested are invited to assist, and will report at the headquarters of the Association at the Lindell Hotel, on the 19th or 20th, where they will be informed of the exact place of meeting. The meetings of the Association will begin on the morning of the 21st of August. Prof. J. K. Rees, at St. Louis, will give information to members about car fares and accommodation.

B. PICKMAN MANN, Secretary.

NOTICE.—In consequence of a series of uncontrollable mishaps, the issue of the present number has been delayed nearly a month beyond its usual time.