

P. 146, gen. 16 and 17 are placed by Reuter in the Bryocorini, 18 and 19 in the Macrolophini.

P. 147, add "Family 3a, Polycetenidæ.

"Genus 1. *Polycetes*, Giglioli, 1864, Q. Journ. Micr. Sci., IV, 25, type *molossus*, Gigl., Pl. Ib., figs. 13-14.

"Genus 2. *Euroctenes*, gen. nov., type *lyrae* (C. O. Waterh., 1879, T. E. S., London, Pl. IX, figs 1-2).

"Genus 3. *Eoctenes*, gen. nov., type *spasmæ* (C. O. W., op. c., figs. 3-4).

"Genus 4. *Hesperoctenes*, gen. nov., type *fumarius* (Westw., 1874, Thesaurus Ent. Oxon., Pl. 38)."

The characters of these four genera have been indicated by various authors, but only one, *Polycetes*, has been named:

1. Palæogæic forms; posterior legs comparatively short and stout, about half the length of the bug; the claws unequal, one nearly simple, the other large and bent, its basal tubercle nearly as long as the simple claw 2.
- 1a. Neogæic forms; posterior legs as long as the bug; claws nearly equal, with a small tooth at the base 1, *Hesperoctenes*, mihi.
2. Antennæ long, the third segment almost (or more than) as long as the fourth 3.
- 2a. Antennæ shorter, third and fourth segments about equally long 4, *Eoctenes*, mihi.
3. Head medianly wider than long. Pronotum transverse 3, *Polycetes*, Gigl.
- 3a. Head medianly longer than wider. Pronotum elongate 2, *Euroctenes*, mihi.

P. 147, delete entry (on p. 147 only) of Family 5. and read "Family 5, Dipsocoridæ (Monogr., as *Ceratocombidæ*, Reuter, 1891, Act. Soc. Sci., Fenn., XIX, No. 6, pp. 1-28, Pl.).

Subfamily 1, Dipsocorinæ.

"Genus 1. *Lichenobia*, Bærensprung, 1857, Berlin Ent. Zeit., I, 165 (= *Ceratocombus*, Fieber, 1860), Wien. Ent. Mon., IV, 267, type *muscorum*, Fall. (= *Coleoptrata*, Zett.), type *ferruginea*, Baer. (= *Coleoptrata*, Zett.), fig., Signoret, 1852, A. S. E., France (2) X. Pl. 16, f. 3, as *Astemma Mulsanti*."

N. B.—*Ceratocombus* was only named, not described, by Signoret.

P. 148, line 3, read "*Trichotonannus*."

P. 148, date of footnote 20 is 1836.