

Section 4. Toes, 4, without web; bill, over 1.10 and less than 1.90.




Tringa alpina pacifica.

Bill, decurved near tip; one or more of inner secondaries, almost entirely white; upper tail coverts, not white, barred with black; legs and feet, black. Spring birds have black on the belly, and back, rufous brown and black. Fall birds have the belly white and back gray.

**Red-backed Sandpiper.
American Dunlin.**

Tringa alpina pacifica.

See page 49.




Tringa ferruginea.
Tail and upper tail coverts.

Bill, decurved near the tip; upper tail coverts, white, banded with black or dark brown.

Curlew Sandpiper.

Tringa ferruginea.

See page 50.



Tringa maculata.
Tail and upper tail coverts.

Bill, nearly straight; back, marked with tawny and black; breast with numerous narrow, brown streaks; none of inner secondaries almost entirely white; lower rump and upper tail coverts, black; the feathers more or less tipped with buff. Two middle tail feathers longer than the others.

**Pectoral Sandpiper
Grass Bird.**

Tringa maculata.

See page 45.



Tringa maculata.

Bill, nearly straight; back, dark; feathers edged with ashy or buff; breast, grayish without brown streaks; one or more of inner secondaries almost entirely white; legs and feet, yellow in life, pale brown in dried skin.

Purple Sandpiper.

Tringa maritima.

See page 44.