

7. *HELCON PEDALIS*. *N. sp.*

♂ ♀.—Same form and sculpture as *fulvipes*, from which it differs by the posterior femora except base, and their tibiae and tarsi being black. Length .40-.48 inch.

Hudson's Bay; Massachusetts.

## MICRO - LEPIDOPTERA.

BY V. T. CHAMBERS, COVINGTON, KENTUCKY.

Continued from Page 50.

## TINEA.

1. *T. eunitariælla*. *N. sp.*

Black; head and face rufous; palpi grayish white; antennae yellowish gray, annulate with black, tips white; wings black, with a costal and dorsal white spot opposite each other just before the middle (sometimes united, forming a fascia), a white fascia (sometimes interrupted) beyond the middle, a costal white spot in the apical portion of the wing, and near the apex an obliquely curved costal white streak; apical portion of the wing bronzy, iridescent, ciliae grayish brown; legs silvery white, in parts tinged with fuscous; posterior wing fuscous. *Alar ex.* less than  $\frac{1}{3}$  of an inch.

The larva is found upon old stone walls and monuments in cemeteries. I do not know whether it feeds upon the hairs contained in the mortar of the walls or upon the mortar itself, or upon Lichens, but upon the wall where I have found it most abundantly, I have never found a trace of Lichens. The case is composed of silk and grains of lime. It is flattened, with the under surface truncate at each end, and the upper surface projects in shape something like the bowl of a spoon at each end; the sides are emarginate near each end. I have lost my notes upon the larva. *Hab.* Kentucky and the Gulf States.

It is one of the handsomest *Tineæ* known to me.

2. *T. Orleansella*. *N. sp.*

Straw color or pale yellowish, thickly dusted with fuscous; a discal fuscous spot about the middle of the wing, and another opposite to it on