

"*Female*. Brown, cinereous beneath. Palpi obliquely ascending; third joint conical, less than one-fourth of the length of the second, abdomen slightly crested, extending beyond the hind wings; tip tawny. Forewings with indistinct undulating pale lines, and with whitish-bordered distinct orbicular and reniform marks; orbicular mark oblique; reniform mark incomplete, bordered, its disk mostly whitish and traversed by an ochraceous streak which extends far beyond it; three whitish costal subapical points. Length of the body 8 lines; of the wings 20 lines."

3. *A. glaucovaria*, Walker M.S.S., N. Sp.

"*Mas*. Cinerea; palpi oblique ascendentes; abdomen cristatum; alæ anticæ cervino subinctæ, glauco notatæ, lineis nigricantibus deviis undulatis, linea submarginali cuneata, lunulis marginalibus guttisque costalibus nigris; posticæ pisco late marginalatæ."

"*Male*. Cinereous. Palpi obliquely ascending; third joint very short, not more than one-sixth of the length of the second. Abdomen crested. Forewings slightly tinged with fawn-colour, marked here and there with glaucous, which hue most prevalent on the large reniform mark; lines blackish, irregular, undulating; submarginal line with uniform angles; marginal lunules and costal dots black; orbicular mark large, oblique, short-oval. Hind-wings with a broad brown border; fringe whitish; underside with the discal mark and the submarginal line brownish. Length of the body 7 lines; of the wings 14 lines." Walker M.S.S.

3. *A. modica*, Guén.—July.

*Distribution*.—New York, (Doubleday).

*Genus* 7. *Miana*, Steph.

4. *M. undulifera*, Walker.—Rare, July.

*Distribution*.—St. John's Bluffs, East Florida, (Doubleday).

*Genus* 8. *Celæna*, Steph.

1. *C?* *contrahens*, Walker M.S.S., N. Sp.—Common, July.

"*Mas*. Pallide cinereo-cervina; abdomen albidum; alæ lunulis marginalibus fuscis; anticæ lineis tribus pallidis nigricante marginatis, 1 a 2 a que undulatis, 3 a flexa, orbiculari et reniformi pallido marginatis; postice albidæ, margine cinereo diffuso."

"*Male*. Pale cinereous fawn-colour. Palpi obliquely ascending; third joint lanceolate, less than half the length of the second.