

- | | | | |
|---|---|--|---------------------|
| 1 | { | Metatarsus I., with one false articulation | <i>californicus</i> |
| | | Metatarsi I., with several false articulations | 2 |
| 2 | { | Dorsum smooth | <i>maculosum</i> |
| | | Dorsum granulate | <i>grande</i> |

Leptobunus grande, Say.

Liobunum similis, Weed.

Va., D. C., Ohio, Ill.

Leptobunus maculosum, Wood.

Pa., W. Va., Ohio.

Leptobunus californicus, n. sp.

Length, 6.6 mm. ; femur I. 2.7 mm.

Colour—white above, mottled with brown and black, the vase-mark indefinite; beneath whitish, with a few brown spots; legs yellowish, with an apical brown ring on each joint. Eye-tubercle low and smooth; body smooth; legs smooth, except the tarsi, which have spines at each articulation. Legs quite stout; metatarsus I. with one and tibia II. with two false articulations; palpi with the fourth joint longer than the third; fifth joint about equal to the third and fourth together.

Southern California (Davidson).

LIOBUNUM, Koch.

Forbesium, Weed.

The genus *Forbesium* is based on young forms of previously described species. *Liobunum* is very rich in species; many forms remain to be described. I will mention but one, readily recognized by having two large yellow spots.

Liobunum bimaculatum, n. sp.

Length, 4.2 mm. ; width of abdomen, 3.5 mm. ; femur I., 11.9 mm. ; femur II., 19.5 mm.

Colour—dark brown, with two large yellow spots over the juncture of the cephalothorax and abdomen; the eye-tubercle black; venter, palpi and legs yellowish; patellæ and tibiæ at tip brownish; metatarsi and tarsi somewhat brownish. Body short and broad; the end of abdomen bent under; the side-pieces of the sternum appear to be completely separated from the sternum proper; coxæ, venter and sternum with a few granules. Eye-tubercle moderately high; more than its length from the anterior margin, and with two rows of spines above. Abdomen with a few transverse rows of small, stiff hairs. Legs extremely long, with some small spines