

2. *Ceratomya*, Chev.

C. caminea, Fabr. By sweeping herbage on the Mountain, June 1857.

3. *Diabrotica*, Chev.

D. vittata, Fabr. Very abundant on melon and cucumber vines, June to October.

4. *Haltica*, Illig.

H. alzi, Harris MSS. In vast numbers on alder-bushes on the common at Laprairie, September.

H. splendida, Dej. Cat. Very abundant on grape-vines, June.

H. collaris, Fabr. Rare.

H. frontalis, Fabr. Abundant in heads of Thistles, July and August.

H. pubescens, Erichs. Very abundant by sweeping grass, June.

H. striolata, Illig. Very abundant, May and June.

5. *Psylliodes*, Latr.

P. punctulata, Mels. By sweeping grass, Logan's farm, June.

6. *Dibolia*, Satr.

D. ærea, Mels. By sweeping herbage on the Mountain, June.

4. Fam. *Chrysomelidæ*, Leach.1. *Heteraspis*, Dej.

H. curtipennis? Mels. Swept from herbage on the Mountain, June.

2. *Bromius*, Cherv.

B. vitis, Fabr. May.

3. *Doryphora*, Fabr. (See Proc. Acad. Nat. Sci. Phila., Vol. VIII. p. 30.)

D. trimaculata, Fabr. Abundant on the milkweed (*Asclepias cornuti*), June and July.

4. *Chrysomela*, Linn. (See Proc. Acad. Nat. Sci. Phila., Vol. VIII. p. 32.)

C. scalaris, Lec. Abundant.

C. Philadelphica, Linn. Abundant on Willows, June and July.

C. Bigsbyana, Kirby. Not so numerous as the last species.

C. (Helodes) trivittata, Say. Very abundant by sweeping grass, June.

C. (Phædon) polygoni, Linn. (*cæruleipennis*, Say.) Very abundant on *Polygonum ariculare* (knot-grass) in waste places.

5. *Chrysochus*, Cherv.

C. auratus, Fabr. Abundant on dogbane (*Apocynum androsemi-folium*), July.

6. *Chalcophana*.

C. picipes. On elm-leaves; not very common. ? (Abundant at Sorel.)

PSEUDOTRIMERA.

1. Fam. *Endomychidæ*, Leach.

(See Proc. Acad. Nat. Sci. Phila., Vol. VI. p. 357.)

1. *Endomychidæ*, Leach.