

27. *ESPERIOPSIS QUATSINOENSIS*, Lambe. 1892.

Described in Transactions, Royal Society of Canada, vol. X, p. 69, pl. iii, figs. 8, 9, and pl. v, figs. 8, 8a, 8b, 8c; types in museum of the Geological Survey of Canada.

*Type localities*—West coast of Vancouver Island, B.C., north of Quatsino Sound, and near Lasqueti Island, Strait of Georgia.

*Distribution*—Behring Sea and North Pacific Ocean as far south as the State of Washington.

28. *ESPERIOPSIS LAXA*, Lambe. 1892.

Described in Transactions, Royal Society of Canada, vol. X, p. 70, pl. iii, fig. 10, and pl. v, figs. 13, 13a, 13b, 13c; type in museum of the Geological Survey of Canada.

*Type locality*—Oyster Bay, Vancouver Island, B.C.

29. *CLADORHIZA ABYSSICOLA*, M. Sars. 1872.

*Distribution*—Between Anticosti and the Gaspé Peninsula, Gulf of St. Lawrence (Whiteaves, 37); coast of New England (Verrill, 33); Baffin Bay (Fristedt, 12).

*Foreign distribution*—Coast of Norway (Sars, 25); between the north coast of Scotland and the Faroe Islands (Carter).

30. *CLADORHIZA GRANDIS*, Verrill. 1879. (? syn. *C. NOBILIS*, Fristedt. 1887.)

*Distribution*—Off the coast of Nova Scotia (Verrill, 32); eastern coast of Greenland (Fristedt).

31. *CLADORHIZA NORDENSKIOLDII*, Fristedt. 1887.

*Locality*—Between Anticosti and the Gaspé Peninsula, Gulf of St. Lawrence (18).

Fristedt's type specimen is from the east coast of Greenland.

32. *CHONDROCLADIA ALASKENSIS*, Lambe. 1894.

Described in Transactions, Royal Society of Canada, vol. XII, p. 119, pl. ii, figs. 7, 7a-e; type material in the U. S. National Museum at Washington, D.C., and in the museum of the Geological Survey of Canada.

*Distribution*—Behring Sea and North Pacific Ocean.

33. *CHONDROCLADIA PULCHRA*, Lambe. 1894.

Described in Transactions, Royal Society of Canada, vol. XII, p. 119, pl. ii, figs. 8, 8a-d; type material in the U. S. National Museum at Washington, D.C., and in the museum of the Geological Survey of Canada.