

Camponotus sp.

Formica sp.

Cremastus sp.

Passalwex mandibularis Cr.

Figites coloradensis Ashm. ♀.

Apanteles sp.

Aspilota pallidipes Ashm.

Pimpla annulipes Brullé.

(8) Horeshoe Bend, 1887—

Urocetus flavicornis Fl.

Dolichoselphus Ashm., is allied to *Atractodes*, with abnormally lengthened maxillary and labial palpi.

LEPIDOPTERA.

Mr. W. H. Edwards has kindly examined many of the butterflies. The two species of *Cidaria* were identified by the Rev. G. D. Hulst.

(1) Lakes of the Clouds, 1887—

Colias meadii Edw.

(2) Near Micawber Mine, 1888—

Nathalis iole Bdv.

(3) Near Brush Creek, June 27, 1889—

Phyciodes camillus Edw.

Nisoniades icelus Lintn.?

Argynnis hesperis Edw.

Lycæna sæpiolus Bdv., ♂, ♀.

“ *oro* Scudd.

(4) Near Micawber Mine, June 27—

Pyrameis cardui L.

“ “ form *minor* Ckll., nov.

Anthocharis ausonides Bdv.

Vanessa milbertii var. *subpallida* Ckll.

Pieris oleracea Bdv.

Gnophaela vermiculata Grote.

Agrotis auxiliaris Grote.

Papilio rutulus Bdv.

Colias scudderii Reak.

Lycæna acmon D. and H.

P. cardui f. *minor* differs from the type only in being much smaller.