

19. *Melitæa phæton*, Drury. Moderately common in June. Pineville.
20. *Phyciodes nycteis*, Doubleday and Hewitson. Moderately common everywhere. Specimens from Lexington, Pineville, Brooklyn Bridge.
21. *Phyciodes tharos*, Drury. Very common all over Kentucky. Seen from May through the season.
22. *Grapta interrogationis*, Fab. One of the most abundant butterflies of Kentucky.
23. *Grapta comma*, Harr. Rather rare. Specimens from Lexington.
24. *Grapta progne*, Cram. Rather rare. Specimens from Clay's Ferry.
25. *Vanessa antiopa*, Linn. Not common, most often seen in the fall.
26. *Pyrameis atalanta*, Linn. Frequently seen throughout the State. Specimens from Lexington.
27. *Pyrameis huntera*, Fab. Rather common. Specimens from Lexington, East Hickman and Bryant.
28. *Pyrameis cardui*, Linn. Common everywhere.
29. *Junonia cænia*, Hubn. Occasional in Eastern Kentucky. Common in western part of State, along the Ohio River. Specimens from Glasgow and Fulton.
30. *Limenitis ursula*, Fab. Frequently seen during the latter part of summer. Specimens from Glasgow, Lexington, Pineville, East Hickman.
31. *Limenitis disippus*, Godt. Not very common. Specimens from Lexington, Glasgow and Tyrone.
32. *Apatura cellis*, Boisd. and Lec. Common everywhere. Specimens from Lexington and Fulton.
33. *Apatura clyton*, Boisd. and Lec. Occasional. Specimens from Lexington, Fulton and Tyrone.
34. *Paphia troglodyta*, Fab. Occasional in Eastern Kentucky. Common locally in Western Kentucky. Found at Lexington, Hopkinsville and Glasgow Junction.
35. *Debis portlandia*, Fab. Moderately common in Western Kentucky. Specimens from Aden Springs and Fulton.
36. *Neonympha gemma*, Hubn. Occasionally seen. Specimens from Fulton.
37. *Neonympha eurytris*, Fab. Rather common in open woods. Specimens from Nortonville, Midland and High Bridge.
38. *Neonympha sosybius*, Fab. Common everywhere in woodland. Specimens from Fulton.