

Bronchitis and Catarrh.

FOR BRONCHITIS.—Take two teaspoonfuls of the cough mixture (that is one teaspoonful of PAIN-KILLER mixed with nine of molasses) three times a day, using as a gargle for the throat, every night and morning, a strong mixture of PAIN-KILLER in water, or milk and water, sweetened with loaf sugar, at the same time bathing the throat in PAIN-KILLER clear. On retiring to bed soak the feet in hot water and apply bottles of hot water to the sides to produce sweating. If the Bowels are costive some gentle physic may be taken.

For evidence in favor of the PAIN-KILLER for Ministers' Sore Throat or Bronchitis, read the following :

GENTS.—The PAIN-KILLER has been a constant occupant of our house for over two years, and a portion of the time it has been the only medicine under our roof. Hardly ever do I have my children complain of being sick, without having them ask in the same sentence for PAIN-KILLER.

For several years before I became acquainted with the PAIN-KILLER, I had suffered a great deal from an affection in my throat, thought by some physicians to be Bronchitis, by others to be what is called Ministers' Sore Throat. At one time it was so severe that I was obliged to give up preaching. Within a few months after I had become acquainted with the PAIN-KILLER, I had another attack from that distressing complaint. I tried my new-found medicine; and, to my astonishment and delight, I found it produced a wonderfully soothing effect. In a short time I was wholly relieved. Since that time, I have had a number of attacks of the same nature, and the PAIN-KILLER has always afforded me relief. About one year since, my wife became subject to severe suffering from Rheumatism; our resort, as usual, was to the PAIN-KILLER, which would always relieve her.

Very truly yours,

EDGAR CADY, OWATONNA, Minn.,
Missionary of the A. B. Home Mission Society.

FOR CATARRH.—The best way to take the PAIN-KILLER for this complaint is to inhale it freely into the Lungs, which can be done by saturating a cloth or sponge with the PAIN-KILLER, and inhaling through it. It is also desirable that the patient should take a teaspoonful of the PAIN-KILLER in half a cup of hot water and sugar, three times a day.