

tative change takes place in this euglobulin, and is associated with the formation of the "anti-body," which is the active agent in the Wassermann reaction. It is here maintained that the two most accurate tests for this euglobulin are the Noguchi butyric acid test and the ammonium sulphate ring test above described. These tests are of the greatest value in distinguishing general paralysis from non-syphilitic affections of the nervous system.

BIBLIOGRAPHY.

Achard, Loeper and Lanbry. "Contribution à la cryoscopie du liquide céphalo-rachidien," *Arch. de méd. expér.*, juillet 1901, p. 567.

Arthus. "Eléments de chimie physiologique," 1900, p. 191.

Babcock. State Hospitals Bulletin, 1896, cited by Schaefer.

Bauer. "Zum Wesen der Wassermann'schen Luesreaktion," *Berl. klin. Woch.*, 1908, S. 834.

Ibid. "Zur Methodik des serologischen Luesnachweises," *Deutsche med. Woch.*, 1908, Nr. 16.

Brandberg. "Jahresber. f. Thierchemie," 1881, Bd. 10, S. 265.

Cimbal. "Chemische, physikalische und morphologische Ergebnisse an 240 Spinalpunktionen und deren diagnostische and therapeutische Verwertung. *Die Therapie der Gegenwart*," Nov., 1906, S. 495.

Coriat. "The Chemical Findings in the Cerebrospinal Fluid and Central Nervous System in various Mental Diseases," *Amer. Journ. of Insanity*, April, 1904, Vol. lx., p. 733.

Decoubaix. "La ponction lombaire dans la psychiatrie," *Bull. de la Soc. de Méd. ment. de Belgique*, 1905, Nr. 119.

Dirksen. "Étude sur la composition chimique et la concentration moléculaire du liquide céphalo-rachidien," *Thèse de Paris*, 1901.