

**Case 22 (CARCINOMA OF THE PANCREAS).**—*Heart*—Large amount of fat, droplets rather large, 3·11 per cent.; iodine value, 122. *Kidney*—1·38 per cent.; iodine value, 125. *Liver*—Moderate amount of fat in lobules and in the carcinomatous areas, 3·21 per cent.; iodine value, 131.

**Case 23 (TUBERCULOSIS).**—*Heart*—Large amount of fat, 2·8 per cent.; iodine value, 118. *Kidney*—Very small amount of fat, 2·41 per cent.; iodine value, 119. *Liver*—Very large amount of fat, 12·81 per cent.; iodine value, 78.

**Case 24 (PANCREATITIS, CHRONIC).**—*Heart*—Very few fat granules, but considerable brown atrophy, 2·09 per cent.; iodine value, 144. *Kidney*—Moderate amount of fatty change, 2·27 per cent.; iodine value, 129. *Liver*—Very large amount of fat, 10·64 per cent.; iodine value, 79.

**Case 25 (TYPHOID WITH ORBITAL CELLULITIS AND MENINGITIS).**—*Heart*—No demonstrable fat, but moderate degree of brown atrophy, 1·77 per cent.; iodine value, 155. *Kidney*—Very small amount of fat present, 1·68 per cent.; iodine value, 141. *Liver*—Large amount of fat, 6·37 per cent.; iodine value, 105.

**Case 26 (PERIPHERAL SEPTICEMIA).**—*Heart*—Small amount of fine droplet fat and some brown atrophy, 1·40 per cent.; iodine value, 141. *Kidney*—Small amount of fatty change, 1·31 per cent.; iodine value, 137. *Liver*—Small amount of fat demonstrable, 2·90 per cent.; iodine value, 133.

**Case 27 (CHRONIC ALCOHOLIC, DEATH FOLLOWING GASTRO-ENTEROSTOMY).** *Heart*—No visible fat, but large amount of brown atrophy, 1·92 per cent.; iodine value, 151. *Kidney*—Small amount of fat, 1·43 per cent.; iodine value, 157. *Liver*—Moderate amount of fat, for the most part in the cells about the periphery of the lobules, 2·95 per cent.; iodine value, 139.

**Case 28 (TUBERCULOSIS).**—*Heart*—Small amount of fat, patchy in its distribution, 2·2 per cent.; iodine value, 127. *Kidney*—Small amount of fat present, 1·6 per cent.; iodine value, 134.

**Case 29 (PNEUMONIA).**—*Heart*—No visible fat, 1·94 per cent.; iodine value, 147. *Kidney*—1·84 per cent.; iodine value, 137. *Liver*—Abundant fat present, 10·31 per cent.; iodine value, 81.

**Case 30 (ADDISON'S DISEASE).**—*Heart*—Very small amount of fat, 2·04 per cent.; iodine value, 121. *Kidney*—Moderate amount of fat, 3·25 per cent.; iodine value, 126. *Liver*—Very abundant fatty change, 11·75 per cent.; iodine value, 83.

**Case 31 (LOBAR PNEUMONIA, MENINGITIS).** *Heart*—No visible fat, but considerable brown atrophy, 1·86 per cent.; iodine value, 152. *Kidney*—Moderate amount of fat, 1·92 per cent.; iodine value, 123. *Liver*—Abundant fat present, 8·31 per cent.; iodine value, 94.

**Case 32 (MILITARY TUBERCULOSIS).**—*Heart*—Moderate amount of fat present, 2·18 per cent.; iodine value, 118. *Liver*—Small amount of demonstrable fat, 3·02 per cent.; iodine value, 131.

**Case 33 (PNEUMONIA).**—*Heart*—No demonstrable fat, some brown atrophy, 1·61 per cent.; iodine value, 159. *Kidney*—No fat present, 1·60 per cent.; iodine value, 143. *Liver*—Small amount of fat present, 3·14 per cent.; iodine value, 140.