

Prothorax dark orange, beset with long fuscous hairs. Meso- and metathorax black or fuscous black. Legs blackish, with many black spines; spurs black.

Fore-wing broad, semi-hyaline, nearly uniformly smoky brown, somewhat suffused with yellowish in basal area; venation fuscous; discal cell very long, twice the length of the first apical cell. Hind-wing similar to the fore-wing, but the basal area not suffused with yellow; colour of the wing lightened in inner marginal area; venation darkish.

Abdomen dark brown, ochraceous toward apex; hind margin of each segment narrowly variegated with dark yellow. In the male the posterior margin of the 9th abdominal segment produced into a quadrangular prolongation in the middle; superior appendage ochraceous, stout, and broad; inferior appendage very long, stout, with the apex furnished with numerous spiny hairs; viewed from above, between the inferior appendages, are seen two slender yellowish processes.

Length of body 5—5.5 mm.; length of fore-wing 9 mm.; length of hind-wing 7 mm.

The type is a single male specimen captured by Mr. S. Yamamura at Minakuchi, Prov. Ohmi, on November 4, 1911.

This species is closely allied to *N. ruficollis*, but can at once be distinguished from the latter by the much smaller size. The structure of the male genitalia and a certain character in wing venation also afford unmistakable distinctive criteria between the two species.

Genus *Moropsyche* Banks.

13. *Moropsyche parvula* Banks.

Moropsyche parvula Banks, Proc. Ent. Soc. Wash., VII, p. 108, pl. III, figs. 3, 8 (1906); Ulmer, Deutch. Ent. Zeit., p. 342 (1908).

This species is unknown to me. I have not yet obtained a specimen, nor have I seen any in Japan.

Habitat—Kiushiu (Hikozan), according to Banks.

Time of appearance—March, according to Banks.