

Also recorded from the Red River (Scudder<sup>1</sup>)\* and Boissevain (Walker).

B. C.—Agassiz. (Walker.)

95. *Xiphidion brevipenne* Scudd.

MAN.—Red River. (Scudder.<sup>1</sup>)

96. *Xiphidion saltans* Scudd.

MAN.—Aug. 8, 21, 1905, 2 ♀'s. (Criddle.)

SASK.—Souris River. (Scudder.<sup>2</sup>)

Both of the Manitoba specimens are brachypterous, the tegmina being shorter than the pronotum.

Subfamily Decticinae.

97. *Neduba carinata* Walk.

B. C.—Wellington. (Caudell.<sup>3</sup>)

98. *Apote notabilis* Scudd.

B. C.—Vancouver Island. (Caudell.<sup>3</sup>)

99. *Anabrus simplex* Haldeman.

MAN.—Aweme, July 15, 1904, 1 ♀ (Willing); Aug. 10, 1903, 1 ♂ (Criddle). Also recorded from Manitoba by Scudder (Psyche, VIII, p. 95, 1897).

SASK.—Radisson, July 29, 1907, 1 ♀ nymph. Also recorded from the Souris River by Scudder.<sup>2</sup>

var. *maculatus* Caudell.

B. C.—Ft. Walsh. (Caudell.<sup>3</sup>)

var. *coloradus* Thomas.

MAN.—Thomas (Bull. U. S. Geol. Surv. Terr., IV, p. 485, 1878).

ALTA.—Macleod. (Caudell.<sup>3</sup>)

The female from Aweme is rather small for the typical form (pronotum 11 mm.), and has faint indications of maculations.

100. *Anabrus longipes* Caudell.

B. C.—Nelson, 2 ♂'s. (Alexander.)

\*Small numerals following an authority's name refer to same numerals in the list of references.