pattern and length of tegnium, though all the species from the Magdalen Islands are brachypterous. One of the females from Bloomfield, a macropterous individual, is strikingly marked, the antennæ, dorsum of head and pronotum being rose-red, the face and lateral lobes of the pronotum grass-green and the tegmina and femora tinged with pinkish.

Mecostethus lineatus (Scudd.)—Bloomfield, 2σ ; Bunbury, 1σ , $1 \circ$; Charlottetown, 1σ (juv.); Tracadie, $1\circ$ (on sphagnum bog); Alright Island, 1σ .

The two males from Bloomfield are less distinctly marked than usual, the pale costal streak of the tegmina being less distinct, the dorsum of the head and the entire pronotum concolorous, without pale lines along the lateral carinæ.

Mecostethus gracilis (Scudd.)—Dundee, 1 ♂.

Camnula peelucida (Scudd.)—Charlottetown, $1 \, \vec{\sigma}$, $2 \, \circ$; Dundee, $1 \, \vec{\sigma}$, $1 \, \circ$; Lake Verde, $1 \, \circ$; Mt. Stewart, $1 \, \vec{\sigma}$; Souris, $1 \, \vec{\sigma}$; Tignish, $1 \, \circ$; West R., Bothwell, $1 \, \vec{\sigma}$; Bic, $10 \, \vec{\sigma}$, $16 \, \circ$; Father Pt., $2 \, \vec{\sigma}$, $2 \, \circ$; Glin House, $1 \, \vec{\sigma}$, $2 \, \circ$; Coffin Island, $1 \, \circ$ (juv.); Grindstone Island, $4 \, \vec{\sigma}$, $1 \, \circ$ ($9 \, \text{iuv}$).

The labels of the Bic specimens bear the following notes: "Dry shaly shore and rocks at head of Ha Ha Bay," "dry shaly fields near Cap Gurgé," "dry rocks" and "dry grassy rocky fields." The specimens from Father Point are labelled "Rocky shores of St. Lawrence River."

Dissosteira carolina (L.)-Point du Chene (2 ♂, juv.)

Circotettix verruculatus (Kirby)—Moncton, 1 ♂; Bic, P. Q., 6 ♂ The Bic specimens are labelled as occurring on "dry exposures," "dry rocky clearing" and "dry grassy limestone exposures."

These specimens are rather pale, resembling *D. carolina*, though one example is rather dark and mottled. The Moncton specimen is almost black

Podisma glacialis glacialis (Scudd.)—Dundee, "east on P.E.I. Ry. toward Douglas," 1 \(\rightarrow \) ("black spruce swamp"). St. Fabien, 1 \(\sigma^7 \) ("bog east of Lac de St. Fabien").

These specimens are quite typical, showing no variation toward the race canadensis E. Walk

Melanoplus fasciatus (Barnston-Walk.)—Dundee, 2 &, 2 &; West River, Bothwell, 1 &; Wolfe Island, 1 &; Alright Island,