

Family PAPILIONIDAE.**Sub-family PIERINÆ.*****Pieris napi*, Esper.**

(PIERIS OLERACEA, Harris.)

A number of specimens have been taken at St. John, Fredericton, and on the Belleisle, but this species is no doubt uncommon throughout the province.

***Pieris rapæ*, Linn.**

This species is an importation from Europe, and is our most common butterfly in New Brunswick. Flying from May to October.

***Collas philodice*, Godt.**

(EURYMUS PHILODICE.)

Very common throughout the province. Flying from May to September, and during the present year as late as October 23rd.

***Collas interior*, Seud.**

(EURYMUS INTERIOR.)

Usually not very common, but during the present year this species was unusually abundant, being much more numerous than *C. philodice*. Flying from July 9th to late in August.

Sub-family PAPILIONINÆ.***Papilio turnus*, Linn.**

(JASONTIDES GLAUCUS, Seud.)

This species is common in every part of New Brunswick, and occasionally very numerous. Flying in June and July.

***Papilio cressphontes*, Cram.**

PAPILIO THOAS, Boisl.

(HERACLIDES CRESSPHONTES.)

Rev. C. J. S. Bethune, in his list of the "Butterflies of the Eastern Provinces of Canada," gives St. John, N. B., as a locality for this species. None of our local collectors have met with it. It is a southern insect and no doubt rarely occurs in this province.

***Papilio brevicauda*, Saunders.**

At present we have no local record of the capture of this species, but Rev. C. J. S. Bethune, in his list of the "Butterflies of the Eastern Provinces of Canada," mentions it as having been taken in Dalhousie, N. B. When the insects of the northern sections of New Brunswick become better known, this butterfly will no doubt be found among them.