

CANADA MEDICAL RECORD

DECEMBER, 1898.

TWO CASES OF TUBAL PREGNANCY—OPERATION—RECOVERY.

By A. LAPHORN SMITH, B.A., M.D., M.R.C.S., Eng.

Fellow of the American Gynecological Society; Fellow of the British Gynecological Society; Gynecologist to the Montreal Dispensary; Professor of Clinical Gynecology in Bishop's University; Surgeon-in-Chief of the Samaritan Hospital for Women; Surgeon to the Western Hospital.

Mrs. B., aet. 33. Came under my care on 5th October. As a girl enjoyed good health. Married at 21. Had three children; last 6 years of age, after which she became a widow. Married again two years ago, since which has been ill. Conception thought to have occurred on three or four occasions, but always aborted at third month with hemorrhage and pains. Had had no menstruation for three months previous to present flow, which came on about a month ago with severe pains. Four days ago she was taken with a slight flooding and such severe pains that she became unconscious, and does not know what came away. Her physician, who had attended her for five weeks, and already diagnosed disease of the ovaries and tubes, now came to the conclusion that there might be a tubal pregnancy, and called me in consultation. I confirmed both diagnoses; that is, the presence of ruptured tubal pregnancy in a diseased tube, and advised her being brought to the Samaritan in order to remove it. She came at once in the ambulance, but her pulse was 110°, and she was vomiting constantly, so I decided to wait until she was in better condition, at the same time