

**Ozena.**—E. J. Moure (*Berl. Klin. Woch.*) applies the nose douche with rhythmic jets as the patient breathes, and the main elements for success are regularity and extreme care. Expulsion of the crusts may be facilitated by vibration massage of the mucous membrane, spraying the nose with a five or twenty per cent. silver nitrate solution. Irregularity of the passage should be corrected with paraffin injections. General treatment should be carried out persistently.

---

**Tobacco Amblyopia.**—De Waele (*Arch. d'Oph.*) treats tobacco amblyopia with .2 gram solutions of lecithine suspended in normal saline solution. There is no local reaction and the pain only lasts a few hours.

---

**Asthma.**—Straübli (*Mun. Med. Woch.*) recommends the use as an inhalation of adrenalin. He has specially devised an apparatus for the inhalations. In severe attacks he advises 20 drops of 1-1000 adrenalin and 2 drops of a solution of atropin and cocaine (sulphate of atropin, 0.1; hydrochlorate of cocaine, 0.25; distilled water, 10.0) to be placed in the apparatus and inhaled.

---

**Uncontrollable Vomiting in Infants.**—Variot, Lavialle and Rousselot (*Bull. de la Soc. de Péd.*), in this report, add fifteen new cases to the twenty they previously reported, thus confirming their treatment of the most tenacious and inveterate vomiting in young infants by sweetened condensed or ordinary milk. There is a prompt sedative action. To the heated milk, about 10 per cent. sugar is added, which sweetens it about the same as condensed milk. This renders the casein more like human milk; and there is some chemical action on the albuminoids from the large amount of sugar under the action of heat.

---

**Acute Colitis in Young Children.**—V. Hatinel and P. Nobécourt (*Arch. de Med. des Enfants*) do not regard infection as the cause of acute colitis, but the result. The aim, in treatment, should be to clean out the stomach and intestines and modify