

of the staminal tube; in our present individual the leaves do not materially differ, nor do we find any other difference from *a.*, except that the bractæa of the involucre are larger and almost foliaceous.

1. *Glaux maritima*? *L.*

A singular state of this plant, as it appears to us, without flowers, with small imperfect leaves, and with the stem singularly inflated at the joints, is in Mr Douglas' collection.

ORD. XLVII. PLANTAGINÆÆ. *Juss.*

1. *Plantago gnaphalioides*. *Nutt. Gen. Am.* 1. p. 100. *Hook. Fl. Bor. Am.* v. 2. p. 124.
—*P. lagopus*. *Ph.* (non *Linn.*).

ORD. XLVIII. NYCTAGINÆÆ. *Juss.*

1. *Abronia mellifera*. *Dougl. in Bot. Mag. t.* 2879. *Hook. Fl. Bor. Am.* v. 2. p. 125.
2. *A. arenaria*. *Menz.—Hook. Ex. Fl. t.* 193. *Fl. Bor. Am.* 2. p. 125.

Found in California first by *Mr Menzies*.

1. *Chenopodium*? *spinosum*. *Hook. Fl. Bor. Am.* v. 2. p. 127.

North California. *Douglas*. 1826.

ORD. XLIX. POLYGONEÆÆ. *Juss.*

ERIOGONUM. *Mich.*

Involucrum tubulosum, campanulatum v. cyathiforme, vix angulatum, subæqualiter 6-dentatum, multiflorum. *Receptaculum* bracteolis intra pedicellos instructum. *Perianthia* exserta, profunde 6-fida. *Benth. in Linn. Trans.* v. 17. p. 407.

§ FASCICULATA. *Benth.*

1. *E. parvifolium* (*Sm. in Rees' Cycl.*); fruticosum, foliis brevissime petiolatis ovatis margine revolutis undulatis subtus lanatis, involucris sublanatis. *Benth. l. c.* p. 411. *Hook. et Arn. supra*, p. 158.

Upper California. *Menzies. Douglas.*

Folia $\frac{1}{2}$ - $\frac{3}{4}$ pollicaria, basi truncata vel subcordata. *Pedunculi* omnes simplices vel terminalis ramosus. *Capitula* pauca lateralia et terminalia, fructifera rubescentia. *Bractea* sub capitulo pauca, parva, ovata vel oblonga. *Involucra* in capitulo plurima, sessilia, tubuloso-campanulata. *Bracteola* intra involucrum plurimæ. *Perianthii laciniæ* inter se subæquales. *Benth.*

2. *E. fasciculatum* (*Benth.*); fruticosum, foliis oblongo-ellipticis linearibusve basi angustatis margine revolutis glabris vel subtus tenuiter albo-tomentosis, involucris glabriusculis. *Benth. l. c.* p. 411.

Upper California. *Menzies. Douglas.*

Folia $\frac{1}{2}$ - $\frac{3}{4}$ -pollicaria, demum glabra. *Pedunculi* longiores quam in *E. parvifolio*, terminalis sæpissime umbellifer, radiis 2-3 inæqualiter elongatis, 1-2 brevissimis vel uno alterove intra radios sessilibus. *Bractea* sub capitulo numerosa, oblongo-lineares. *Bracteola* intra involucra minutissimæ. *Involucra* et *flores E. parvifolii*. *Benth.*