

D. versicolorella. N. sp.

Head and palpi ochreous, thickly dusted with brown; a brown annulus around the base of the third joint of the palpi, and another before the apex. Antennae dark brown, faintly annulate with ochreous, and with five or six white annulations in the apical portion. Thorax and anterior wings ochreous, thickly dusted with dark brown; a little less thickly in the basal fourth of the wing, with a brown streak across the base of the wing, and a brown streak extending obliquely from the costa about the basal fourth, to the fold, which, however, is scarcely distinguishable from the thickly dusted portion of the wing behind it; no brown spot at the apex. Posterior wings pale fuscous: abdomen ochreous, the segments not margined with whitish, as in the preceding species. Alar ex. $\frac{1}{2}$ of an inch. Captured in Kentucky in May.

D. bicosto-maculella. N. sp.

Head pale yellowish, the vertex dusted with fuscous; antennae dark brown: second joint of the palpi pale yellowish, tipped with brown beneath: third joint brown, sprinkled above with pale yellowish: thorax and anterior wings blackish, or very dark brown, with ochreous and gray intermixed, with a small and indistinct ochreous spot on the costa, near the base, and another distinct costalous at the beginning of the ciliae, and an opposite dorsal one; ciliae yellowish-ochreous. There are several rather undefined irregular blackish spots or patches on the wings, which, to the naked eye, appear to form three irregular transverse bands, not very definite in outline, one of which adjoins each of the costal ochreous spots, whilst the other is between them. Alar ex. $\frac{1}{2}$ inch. Kentucky.

D. querciella. N. sp.

This species is a *Depressaria* in every respect except that there is a small but very distinct tuft of erect scales at the apex of the thorax. I have but a single specimen, which, however, is in perfect condition, and shows no sign of any injury, so that I cannot doubt that the tuft is a normal structure.

Antennae dark brown or rather blackish, annulate with white; palpi iron gray; head silvery, flecked with dark brown or blackish scales; thorax iron gray, the tuft being ochreous; anterior wings dark iron gray, with a distinct small blackish spot on the costa at about the basal fourth, and two other smaller ones on the costa, one about the middle, and the other at the beginning of the apical ciliae; there are three or four similar small ones on the disc: ciliae ochreous; posterior wings pale