

Date.	Sp. Gr.	Drink taken.	Urine	Sugar for 24 hours.	Urea for 24 hours.	Aceto Acetic Reaction.	MEDICINE.	SYMPTOMS.
1886 Dec.	60 F.	oz.	oz.	oz.				
1	1.028	95	100	3.846	1.752	None.	1 m of 1% nitroglycerine and 1 gr. co deia.	
2	1.0278	60	70	2.692	1.423	"	" " " "	Cramp during night.
3	1.024	43	46	0.46	1.288	"	" " " "	"
4	1.035	50	53	0.60	1.171	"	" " " "	"
5	1.0195	50	61	1.053	0.966	Strong.	" " " "	No sleep and pain inside and back.
6	1.023	44	56	1.166	0.9367	"	" " " "	"
7	1.0245	57	60	1.156	1.212	"	" " " "	"
8	1.0195	58	60	0.577	1.003	Still m're	" with 15 grs. am. cl.	Pain worse.
9	1.028	54	40	1.000	0.862	"	" " " "	"
10	1.026	64	63	2.423	1.053	"	" with 20grs. Na.Br.	"
11	1.027	80	90	2.314	1.212	Strong.	" " " "	Pain less, cold perspiration.
12	1.0288	69	81	3.374	1.288	Faint.	" " " "	"
13	1.029	53	70	2.916	0.528	"	" " " "	Pain gone.
14	1.029	85	90	3.749	1.212	None.	" " " "	"
15	1.02	70	100	5.55	0.970	"	" " " "	"
16	1.026	80	108	4.914	1.10	"	" " " "	Sleep good.
17	1.0285	46	58	3.219	0.86	Faint.	1m of 1% nitro-glycer. t.d. & a little milk	"
18	1.031	60	90	4.995	1.115	"	1m and two half minims	Pain in back again.
19	1.031	66	88	4.884	0.973	"	" + iron & chloric ether	Pain worse.
20	1.035	50	56	2.800	0.906	"	" " " "	"
21	1.032	56	80	4.208	1.167	"	" " " "	"
22	1.028	39	54	1.645	1.104	Stronger.	" " " "	Pain better.
23	1.033	50	52	2.6	0.803	"	3 x 1/2 m doses nitro-glycer.	No pain; weak.
24	1.032	54	70	3.5	1.094	"	" " " "	"
25	1.031	40	62	2.384	0.819	"	2x 1/2 m doses & 1m dose. Biscuit & ale	"
26	1.031	90	144	7.992	1.164	None.	" " " "	"
27	1.031	60	82	4.10	0.973	"	" " " "	"
28	1.033	40	56	2.667	0.996	Faint.	" " Diet strict.	"
29	1.034	58	72	3.6	1.307	"	" " " "	"
30	1.0335	43	60	2.730	0.970	Stronger.	" " " "	"
31	1.032	60	80	3.809	1.161	"	" " " "	Feels stronger.
				92.926				

Note rise of sugar on 26th after starch food and ale.

1887 Jan.								
1	1.035	70	80	4.00	1.208	Faint.	No medicine.	Diet not strict.
2	1.033	25	30	0.872	0.728	None.	" " " "	"
3	1.032	75	102	2.623	1.100	"	" " " "	"
4	1.035	56	82	4.313	1.061	Faint.	" " " "	Cramp at night.
5	1.034	56	80	4.44	1.078	"	2m nitro-glycerine p. day	Acid vomiting all night.
6	1.038	54	50	2.50	0.943	"	" " " "	Legs heavy and tired.
7	1.033	43	52	2.261	0.953	Strong.	" " " "	"
8	1.031	62	70	3.50	0.800	Faint.	" " " "	"
9	1.032	64	94	4.997	0.861	None.	" " " "	"
10	1.033	62	85	4.25	0.908	"	" " " "	"
11	1.032	68	90	4.995	1.164	"	" " " "	"
12	1.035	38	42	1.534	0.974	"	" " " "	"
13	1.0315	36	42	1.164	0.906	Faint.	" " " "	"
14	1.032	42	54	1.690	1.193	None.	" " " "	cold sweat.
15	1.0315	48	78	3.549	1.323	"	" " " "	"
16	1.035	50	70	3.885	1.132	"	" " " "	"
17	1.036	48	60	3.996	0.936	"	" " " "	"
18	1.034	31	66	3.472	1.067	Faint.	" " " "	"
19	1.036	46	60	3.528	0.792	"	Gluten and strict diet	Better nights.
20	1.035	74	75	3.085	1.011	Strong.	" " " "	"
21	1.033	56	60	3.156	0.905	"	" " " "	"
22	1.033	64	76	3.8	1.515	Faint.	" " " "	"
23	1.031	98	101	5.05	1.307	None.	" " " "	"
24	1.032	36	47	2.238	0.766	"	None.	"
25	1.0355	54	66	3.880	1.067	Faint.	" " " "	"
26	1.0315	62	66	2.749	1.103	"	" " " "	"
27	1.0328	54	60	2.857	1.03	"	" " " "	"
28	1.0332	64	68	3.40	1.45	"	A 1m dose	"
29	1.035	49	65	3.6075	1.052	"	Two 1m doses.	Vision slightly affected.
30	1.037	51	60	2.857	1.07	"	" " " "	"
31	1.0325	73	76	3.8	1.25	"	" " " "	"
				102.082				