

- Cardiophorus gagates*, Er.—On oak leaves (May).
Elater manipularis, Cand.—Under decaying linden bark (Nov.)
Elater nigricollis, Hbst.—In hickory stump (Oct.) ; and poplar stump (Dec.).
Elater linteus, Say.—In decaying poplar stump (fall).
Elater apicatus, Say.—In decaying poplar stump (Dec.)
Agriotes oblongicollis, Melsh.—On leaves of hazel (May) ; and beech (May, June).
Limonius auripilis, Say.—On beech foliage (May, June).
Limonius basillaris, Say.—On new oak and beech growth (May, June).
Acmæodera culta, Web.—On cut oak wood (July).
Agilus ruficollis, F., *otiosus*, Say, *bilineatus*, Web.—On cut oak wood and rails (July).
Brachys ovata, Web.—On foliage of beech and oak (May to July).
Brachys acrota, Melsh.—Eating oak and hazel leaves (May) ; leaves of beech, elm, ironwood (*Carpinus*), (May, June).
Brachys aeruginosa, Gory.—On leaves of beech, elm, ironwood (*Carpinus*), (May, June).
Thanasimus dubius, F.—In numbers on piles of oak wood (May). In coitu (May 20).
Thaneroclerus sanguineus, Say.—Under bark oak stumps (Oct.) ; under bark butternut log (May).
Dichelonycha elongata, Fab.—On leaves of oak (May).
Lachnosterna prunina, Lec.—On foliage of red oak (May, June).
Lachnosterna crenulata, Fröh.—On foliage of red oak (May).
Lachnosterna tristis, F.—Swarming in red oaks (May).
Anomala undulata, Melsh.—Swarming in red oaks (May).
Strigoderma arboricola, F.—On red oaks (May). The last five species occur mostly in the evening.
Cotalpa lanigera, L.—Eating leaves of dwarf willows (*Salix cordata*), (May, June).
Parandra brunnea, F.—Under bark of linden and butternut logs.
Smodicum cucujiforme, Say.—Many specimens found on a white oak fence post, beside small holes in post from which they had emerged (June, July).