


profuse perspiration; four hours after the first dose, urine and liquid fæces were passed freely in bed, and the urine



continued to pass away in large quantity during the night. Six hours after the quinine the temperature had fallen to 103°, pulse 112, and fourteen hours after quinine to 100.4°, pulse 90. Dr. Drake saw him the following day, and ordered quin., gr. v, to be given every four hours; four such doses were given when he was allowed up. All bad symptoms disappeared, and he was discharged from Hospital on the morning of July 17 having been under treatment for three days and a half, and having taken altogether fifty grains of quinine in divided doses.

The quinine was administered by the mouth in the form of 10 grain cachets: if it had been impossible to give the quinine in this manner, it would have been administered hypodermically.

The accompanying chart shews the rapid fall of the temperature under the quinine treatment.

PROCEEDINGS OF THE ANNUAL MEETING OF THE CANADIAN MEDICAL ASSOCIATION.

The ninth annual meeting of the Canadian Medical Association was held in the City of Toronto on the second and third days of August, ult:

The morning session was occupied with routine business, and the proceedings in the afternoon commenced with the address by the President, Dr. Hodder, of which the following is a synopsis.—“Gentlemen, we meet together this [morning, to celebrate the ninth annual meeting of the Canadian Medical Association, and from the large number of visitors and members whom I see