THE BEST ANTISF

Both Internal and

LISTEX

Antiseptic, Prophylactic, Deodorant, Non-Toxic, Non-Irritant, Non-Escharotic, Absolutely Safe, Agreeable, Scientific and Strictly Professional.

FORMIULA.—Listerine is the essential antiseptic constituent of Thyme, Encalyptus, Eaptisia, Gaultheria and Mentha Arcensis, in combination.

Each fluid drachm also contains two grains of refined and purified Benzo-boracic Acid.

DOSE.—Internally: One teaspoonful three or more times a day (as indicated), either full strength or diluted, as necessary for varied conditions.

LESTEUINE is a well-proven antiseptic agent—an antizymotic—especially adapted to intermed use, and to make and maintain surgical cleanliness—asepsis—in the treatment of all parts of the human body, whether by spray, interaction, atomization, or simple local application, and therefore characterized by its particular adaptibility to the field of

PREVENTIVE MEDICINE INDIVIDUAL PROPHYLAXIS.

Physicians interested in LISTERINE will please send us their address, and receive by return mail our new and complete pamphlet of

Thysicians interested in LISTERINE will piece of quarto pages, embodying:

A TABULATED EXHIBIT of the action of LISTERINE upon inert laboratory compounds.

A TABULATED EXHAUSTIVE DEPORTS and Clinical observations from all sources, confirming the utility of LISTERINE as a general antiseptic for both internal and external use: and particularly MICROSCOPIC OBSERVATIONS, showing the comparative value and availability of various antiseptics in the treatment of diseases of the oral cavity, by W. D. Miller, A. B., Ph. D., D. D. S., Professor of Operative and Clinical Dentistry, University of Berlin, from whose deductions LISTERINE appears to be the most acceptable prophylactic for the care and preservation of the teeth.

Diseases of the Uric Acid Diathesis.

LAMBERTS

Lithiated Hudrangea.

KIDNEY ALTERATIVE-ANTI-LITHIC.

FORNULA.—Each fluid drachm of "Lithiated Hydrangea" represents thirty grains of fresh Hydrangea and three grains of chemically pure Benzo-Salicylate of Lithia. Prepared by our improved process of osmosis, it is invariably of definite and uniform therapeutic strength, and Benzo-Salicylate of Lithia. Prepared by our hence can be depended upon in clinical practice. DOSE.—One or two teaspoonfuls four times a day (preferably between meals).

Urinary Calculus, Gout, Rheumatism, Bright's Disease. Diabetes, Cystitis, Hæmaturia, Albuminuria and Vesical Irritations generally.

We have had prepared for the convenience of physicians DIETETIC NOTES (sample of which is herewith shown), suggesting the articles of food to be allowed or prohibited in several of these diseases

A neatly bound book of these DIETETIC NOTES, each note perforated for the convenience of pysicians in detaching and distributing to their patients, mailed gratis upon request, together with the latest compilation of case reports and clinical observations, bearing upon the treatment

LAMBERT PHARMACAL

St. Louis, Mo., U. S. A.

GOUT

DIETETIC NOTE .- A mixed diet should be adopted, the nitrogenous and saccharine articles being used in limited amounts.

Allowed .- Cooked fruits without much sugar, tea and coffee in moderation Alcoholic stimulants, if used at all, should be in the form of light wines or spirits well diluted. The free ingestion of pure water is important.

Avoid.—Pastry; malt liquors and sweet wines are veritable poisons to these patients.

The Retail Drug Trade promptly supplied with our products by any Wholesale Druggist of Canada, or from our Canadian Depot at Toronto, by W. LLOYD WOOD, Agent,