

**IGHT
LE**

vent.

11, 1869.
Lungs, and
two doses
the bleeding
all signs of
health fully

ELAND.

up,

ES.

ntly fatal
distressing
free from


es. The
soreness,
round the
follow. If
re accom-
also apply
inflamed
ful every
dy Relief
ation to
thicken-
es of the
patient to
. Never
where the
d. This
ing with

l to the
owsiness,
r, sweet-
mptoms.
dose of
and in a
deposits.
rseness,
lis—in a

CROUP.—Bathe the chest and throat with the Ready Relief—give from a tea to a tablespoonful of the Syrup of Ipecac; if this does not vomit in ten minutes, repeat the dose—this will cure Croup—every time. Thousands of children have been saved by this treatment. As the disease is worse in the middle of the night, every one should keep the R. R. Relief and Ipecac on hand. If on the first signs of Croup half a teaspoonful of Resolvent is given the child—if he be 5 or 10 years old—three or four times per day—and the Relief rubbed on the chest and throat—no further trouble will occur.

CONGESTION OF THE LUNGS.—If suddenly seized with Congestion or Inflammation of the Lungs, Pneumonia, or Pleurisy, rub the chest and side, back and spine, freely with Radway's Ready Relief; give the Resolvent in doses of a teaspoonful *every hour*, and take 6 or 8 of Radway's Pills. In the most distressing forms, the patient will breathe free and easy; in a few minutes after the rubbing with the Relief, and swallowing the Resolvent, all danger will be passed as soon as the Pills operate; but it is well to continue the Resolvent and Pills in small doses for several days, and to wear around the body flannel soaked with Ready Relief. If the simple treatment in the above forms of epidemic diseases is followed, life will be preserved.

A VIEW OF THE URINARY ORGANS



No. 1.—The Kidneys. No. 2.—The Ureters. No. 3.—The Bladder. No. 4.—The Rectum.
No. 5.—The Abdominal Aorta. No. 6.—The Renal Capsules.