

feeds upon the Phylloxera and destroys it, associated with the lice and busy in its useful mission.

Since this insect is now known to be at work in Ontario, and probably to a greater extent than we are at present aware of, a condensed account of its life history will probably be interesting to our readers. The figures are from Prof. Riley's excellent reports, and the facts given mainly gleaned from the writings of this and other authors.

Its progress in Europe has been most alarming, inflicting untold losses in the wine making districts. The destruction it has occasioned in France has been so great that it has become a national calamity which the Government has appointed special agents to enquire into; large sums of money have also been offered as prizes to be given to any one who shall discover an efficient remedy for this insect pest. At the same time it has made alarming progress in Portugal, also in Switzerland and some parts of Germany, and among vines under glass in England. It is a native of America, from whence it has doubtless been carried to France; it is common throughout the greater portion of the United States, and in one of its forms in Canada, but our native grape vines seem to endure the attacks of the insect much better than do those of Europe. Recently it has appeared on the Pacific slope in the fertile vineyards of California, where the European varieties are largely cultivated, and hence its introduction there will probably prove disastrous to grape culture.

This insect is found in two different forms: in one instance on the leaf, where it produces greenish red or yellow galls of various shapes and sizes, and is known as the type *Gallaeicola*, or gall-inhabiting; in the other and more destructive form, on the root, known as the type *Radeticola*, or root-inhabiting, causing at first swellings on the young rootlets, followed by decay, which gradually extends to the larger roots as the insects congregate upon them. These two forms will for convenience be treated together.

The first reference made to the gall-producing form was by Dr. Fitch in 1854, in the Transactions of the New York State Agricultural Society, where he described it under the name of *Pemphigus vitifoliae*. Early in June there appear upon the vine leaves small globular or cup-shaped galls of varying sizes; a section of one of these is shown at *d*, figure 15; they are of a greenish red or yellow color, with their outer surface somewhat uneven and woolly. Figure 14 represents a leaf badly infested with these galls. On opening one of the freshly formed galls, it will be found to