

tion, and, while not, in themselves, directly dangerous, may become so by inducing emphysema and bronchiectasis.

In the great majority of chronic and recurrent winter coughs, the basic trouble lies in a low form of inflammation of the bronchial mucous membrane, especially that of the bronchioles.

In many cases I have used Codeia, but lately I have been having much more success with another derivative of opium, *i.e.*, Heroin. In comparing the results obtained from the use of these two drugs, I notice that heroin will not constipate the patient, nor will it have the stupefying effect characteristic of codeine. Another advantage possessed by heroin is that it is effective in young children, in very small doses.

I had been accustomed to prescribe heroin alone, but, about a year ago my attention was called to a preparation of that drug—Glyco-Heroin (Smith). Upon giving it a good trial I found that it gave me better results than obtained when heroin alone was given, and much more quickly. Glyco-Heroin (Smith) has one distinct advantage over plain heroin in that it can be given for a long time without ill effects, and in the class of patients in question this is, indeed, a most important feature. During the past year and a half I have treated a number of cases and recurrent winter coughs with Glyco-Heroin (Smith) and have obtained uniformly good results.

*Example.*—A. L. Salesman. Aged 28. I saw this patient early in the spring of 1903. He is robust and of good habits. He consulted me concerning a constant cough which had troubled him for over a year. It was usually worse in the morning and after meals, and accompanied by expectoration of thick mucopurulent matter. Sometimes blood-stained, and especially so after a severe paroxysm. This circumstance preyed upon his mind considerably—he thought he had consumption. I learned that he had had a severe attack of acute bronchitis during the spring of 1902 and had been coughing ever since. Physical examination excluded tuberculosis. The diagnosis was chronic bronchitis, sequential to acute. The patient was immediately put on Glyco-Heroin (Smith), and the same hygienic measures ordered as in Case 1. Here again the financial condition of the patient precluded change of climate. In addition to the Glyco-Heroin (Smith) the patient was given syrup of hypophosphites as a tonic. I did not see him again until last October. He then reported himself absolutely free from cough. He continued taking the Glyco-Heroin (Smith), and, during the present winter, has not experienced any return of the trouble. In this case a