

The Canadian Entomologist.

VOL. XXXVI.

LONDON, FEBRUARY, 1904.

No. 2

NEW NOCTUIDÆ FROM BRITISH COLUMBIA.

BY HARRISON G. DYAR, U. S. NATIONAL MUSEUM, WASHINGTON, D. C.

Apatela mæsta, n. var.

A very distinct form of the European *leporina*, Linn.; dark gray, fully as dark as *Canadensis*, Smith & Dyar, which it much resembles; but the basal line is broken and the transverse posterior line is dentate as in *leporina*.

Described from 6 specimens, Kaslo, B. C.

U. S. National Museum, type No. 7322.

Apatela griseor, n. var.

A western variety of *innotata*, Guen., slightly larger, the white ground colour more distinctly strewn with black scales and the black marks bordering the usual lines sharper and more pronounced, especially noticeable in the transverse anterior line. In *innotata* this line is almost lost; in *griseor* it is often nearly as obvious as the transverse posterior line.

Described from 16 specimens, Kaslo, B. C.

U. S. National Museum, type No. 7323.

Apatela dolorosa, n. var.

A western form of *distans*, Grote; darker than its eastern ally, both in the tone of the gray ground and the amount of black shadings. It closely resembles *emaculata*, Smith, from the same region, but is a less smoothly gray and has more marked contrasts between the light and dark shades; the transverse posterior line is more distinctly dentate, and there are other slight differences, so that I think we have to do with distinct, although closely allied species.

Described from 3 specimens, Kaslo, B. C.

U. S. National Museum, type No. 7324.

Platyperigea anotha, n. sp.

Luteous, strewn with dark scales. Head and palpi black-brown, thorax and abdomen luteous grayish. Fore wings shining luteous, basal line rather distinct, broken, the costal half supplemented by an outwardly placed shade on inner margin; t.-a. line distinct, single, brown-black,