

The following clinical notes will afford a general idea of its action in a variety of cases :

CASE 1.—Mrs. ——— came to me presenting the following symptoms incident to a delayed menstruation: Persistent headache of a neuralgic character; dull, aching pain in limbs and lumbar region; cramp-like pains in abdomen, and considerable nausea. The menstrual period was overdue seven days, but as yet there was no appearance of flow. Her periods had always been occasions of intense suffering, but had never before been delayed. I began the use of Ergoapiol (Smith), with some misgiving owing to the irritable condition of the stomach. One capsule every three hours was administered without any aggravation of the gastric distress. In twenty hours a normal menstruation was well under way; the flow was slightly increased over that observed on former occasions. The pains had subsided. Ergoapiol (Smith) was administered, one capsule three times a day, during the menstrual period, which terminated in five days. The patient was instructed to return for a quantity of the remedy several days before the next menstrual period. She did so, and, following directions, took one capsule three times a day for three days before expected menstruation. She subsequently reported that during the period—lasting five days—there had been practically no pain, and that the amount of flow was, as far as she could judge, normal.

CASE 2.—Miss ———, aged 30, had been a sufferer for years with dysmenorrhea. For about three years had suffered with leucorrhœa, particularly annoying after each menstrual period. Had undergone treatment at different times for the leucorrhœa and dysmenorrhea, but had never experienced permanent benefit. She had been obliged to spend the couple of days of each period in bed. She consulted me about one week before her period. Examination revealed a purulent discharge oozing from os cervix and a rather large uterus. There was no displacement. She was put upon Ergoapiol (Smith), one capsule three times a day. The onset occurred one day earlier than was expected and was attended with considerable pain. The patient was, however, able to attend to her usual duties, a state of affairs such as had not been experienced for some years. At the onset of the flow Ergoapiol (Smith) was administered, one capsule every two hours. The effect was astonishing. In eight hours the pains had well-nigh subsided and there was practically no discomfort, except some pain in back.

CASE 3.—Miss ———, aged 21, had suffered for two years with irregular and painful menstruation. Had commenced to menstruate when 16, menses being very scanty, but regular and accompanied with but slight degree of suffering. Was never of a very robust physique, but in the main healthy. When