

Under side of hind wings is entirely of bronze colour, somewhat lustrous, excepting at the top portion of each interspace, where there is a whitish spot, and inside of that, one-sixteenth inch from margin, a semicircle of brick colour, very prominent. The dentations of the wing bear the white linear border appearing on upper side.

The specimens in my possession were taken some two hundred miles north from Cochabamba. In general appearance, it resembles closely *Papilio Numitor*, Cram., and that group, and may be a climatic variation of it, but in *Numitor* the greenish-white dashes on hind wing follow more closely the contour of the hind margin, while in this these dashes are more in line with a line drawn from apex to anal angle. My specimens are invariable, showing no difference in the suffusion or size of dashes.

*Dynamine albidula*, sp. nov.

Habitat: Bolivia, near Sicastica. Expanse, 1.25 inches.

Head, palpi, thorax and abdomen above, blackish with gray hairs; beneath, nearly white. Antennæ, black with white annulations at the base of each joint. Club, tipped with tawny. Legs, white.

The lower portion of the fore wing is white, from a line drawn from a point close to the base on inner margin, and extending upwards to and along the median nervure to end of discoidal space, then curving downwards to lower angle, meeting inner margin one-sixteenth inch from angle. The rest of the wing is black with white spots. The discoidal space is heavily dusted with lustrous greenish-blue scales. In the centre of the space is a small white spot. Midway between apex and base is a large white spot, extending from costa towards hind margin, and directly over the summit of the lower white area. Just within hind margin, one-third distance from apex to lower angle, is a smaller white spot, and there is another below it, one-third distance from lower angle to apex. The latter of these merges into the white area of the lower portion of the wing.

Upper side of lower wing is entirely white, excepting a small area of black at the very base, and a narrow border of black along hind margin. This border is black at the upper angle, and turns to grayish at the lower half, disappearing entirely just before it reaches anal angle. Outside of this, on the edge of the margin, is a white thread.

The white area of upper side of fore wing is duplicated on under side. The black of the upper side gives way to a great extent to tawny. The discoidal space is jet black at its upper portion, tawny at basal portion,