Murawjew considers it as the most efficacious, indeed a specific renedy in Favus.

In toothache, as a result of Caries; in this case conjum gives but temporary relief-" in all cases however of a year!" In neuralgic affections the external application of the alkaloid is attended with marked reliefespecially in pairful syphilitic affections of the bones, in which it is given, of merely as a palliative, but as a radical cure. If is found of remarkable service in synovitis, especially when chronic, whether it has a traumatic, catarrhal of scrofulous origin. As an application to wounds, eshecially gun shot wounds, and in that ordema or rheumatic pains, which follow, a salve of conturn proves it of marked service. A few experiments seemed to prove commit to be of service, in scrofulous, and rheumatic inflammation of the eye, inssmuch as it soon dissipated the intolerance of light, the flowing of tears, &c., &c. In scirrhus and cancer, its action seemed to be merely pulliative. In a case of cancer of the bp in a female peasant of 38 years of age, on whom he had operated, the common had been applied diligently for 3 weeks, in form of dressing. In this case all Marawaew's attention was required to guard against danger from inhalation of common vapour, which produced intoxication. In general, it is necessary, that the quality of the alkaloid should not vary, and for this reason he recommends the following: As an application in skin disease, Murawjew uses to one onnce of carron oil (lime water and has ed oil.) or spermaceti ointment 12 to 24 drops of cenera. He rubs the part previously with flannel, then smears the salve, and applies a piece of oil cloth, glove leather, or post paper, and over this a bandage. When it is necessary (as for instance in neuralgia) to apply pure coneia, the skin is first rubbed with spirit of wine or Cologne water, then three or four drops are allowed to fall upon the part, which is then immediate y covered with oil cloth, and a bandage. The salve must be always kept m a warm place and m a well closed vessel (a stoppered bottle is preferable). As a collyrium Murawjew uses 1 to 3 drops to 6 drachms of water and 2 of mucilage; as a clyster 1 to 3 drops in a decostion of starch or Inseed.—Medizinische Neuigkeiten.

## FRENCH.

Injonctions ud. cs dans le traitement de la dyssenteure chronique (Relioux).—Voici le résultat de quelques expériences sur l'emploi des li vements iodés dans le traitement de la dyssenteure qui sont très dignes de l'attention des médecins.

La dyssenterie, dit M. Pelioux, a peur caractères anatomiques incontestables des lésions spéciales de la membrane muqueuse cœco-colique. Dans la forme chronique surtout, qui a fait particulièrement l'objet de mes observations, cette membrane est rouge livide, boursouffée, ou même frappée d'une véritable hypertrophie qui envalut également la tunique musculeuse et le tissu musculaire interposé; la surface interne de l'intestin est parsemée d'ulcères taillés à pic, souvent recouverts d'une exsudation pseudo-membraneuse u de plaques gangreneuses; entre les