

## MEDICINES WHICH STIMULATE THE LIVER.

Podophyllin in small doses, is a stimulant of the liver. During the increased secretion of bile, the percentage amount of special bile solids is not diminished. If the dose be too large, the secretion of bile is not increased. It is a powerful intestinal irritant.

Euonymin is a powerful hepatic stimulant. It is not nearly so powerful an irritant of the intestine as podophyllin.

Sanguinarian is a powerful hepatic stimulant. It also stimulates the intestine, but not nearly so powerfully as podophyllin.

Irisin is a powerful hepatic stimulant. It also stimulates the intestine, but not so powerfully as podophyllin.

Leptandrin is a hepatic stimulant of moderate power. It is a feeble intestinal stimulant.

Colocynth is a powerful hepatic as well as intestinal stimulant. It renders the bile more watery, but increases the secretion of biliary matter.

Jalap is a powerful hepatic as well as intestinal stimulant.

Menispermmin does not stimulate the liver. It slightly irritates the intestinal glands.

Babtinin is a hepatic, and also an intestinal stimulant of considerable power.

Phytolacin is a hepatic stimulant of considerable power. It also slightly stimulates the intestinal glands.

Hydrastin is a moderately powerful hepatic stimulant, and a feeble intestinal stimulant.

Juglandin is a moderately powerful hepatic and mild intestinal stimulant.

Chloride of ammonium is credited with chologogue properties, but it is questionable; nevertheless it certainly stimulates the intestinal glands.

Calomel is a powerful purgative, but whether it stimulates the liver is still *sub judice*.

Corrosive sublimate is a potent hepatic stimulant, but acts feebly on the intestines.

Sulphate of potash is a powerful intestinal irritant, but its action on the liver is variable and unreliable.

Taraxacum is a feeble hepatic stimulant.

Dilute nitro-muriatic acid has a moderate stimulant action on the liver.

Boldo, bromide of potassium, nitrite of potash, and hard soap have each some stimulant action on the liver.

## THE TREATMENT OF EPISTAXIS.

Introduce into the nostril, to a considerable distance upward, a piece of fine sponge, cut to the size and shape necessary to enable it to enter without difficulty, previously soaked in lemon juice or vinegar and water. The patient is to be kept lying on the face for a length of time, with the sponge in place. This, says *Lyon Medical*, is the procedure employed by M. Siredey for controlling epistaxis in typhoid fever patients.—*N. Y. Medical Journal*.

## HOW TO SHRINK HYPERTROPHIED TONSILS BY CAUSTIC APPLICATIONS.

Among various caustics for local use in causing shrinkage of tonsillar hypertrophies, Dr. Chisholm (*Virginia Medical Monthly*) has found the chloride of zinc the most available and the least annoying to the patient. He employs it in the following manner: A wire the size of a fine knitting needle, is roughened for a half inch from one end so that it may hold a fibre of absorbent cotton twisted upon it. Dip this into a saturated solution of chloride of zinc and thrust it to the very bottom of the crypt, and keep it there for several seconds. When withdrawn the whitened orifice marks the cauterization. By renewing the cotton for each follicle several may be thoroughly cauterized at the same sitting without causing any annoying irritation to the throat. A very few applications will cause the gland to shrink, as will be seen one week after the destructive cauterization has been made to the interior of the follicles.—*Medical Record*.

## CARBOLIC ACID IN INDIGESTION.

Berdoo has frequently treated acid dyspepsia with small doses of carbolie acid. He uses a solution suggested by Dr. Fenwick, containing one part of the crystallized acid in four parts of glycerin, and gives from five to ten minims as a dose, either merely diluted or mixed with nux vomica or liquor opii sedativus, the latter being added in case of pain. He does not attempt to explain the action of the drug, but suggests that its efficacy may be due either to its anti-fermentative power or to the anæsthesia which it induces in the gastric mucous membrane.—*N. Y. Medical Journal*.

## IODIDE OF POTASSIUM IN INFLAMED BREAST.

Dr. Samuel Welch thus writes in the *Med. Press*, April 22d: Having been frequently disappointed with the ordinary remedies in the highly troublesome condition arising from the presence of milk in the breast after the death of the child, or in cases of still-born children, and having found that the effects of belladonna are often uncertain, and that purgatives, although certainly useful, are frequently unreliable, I determined to try the effect of iodide of potassium applied locally in the form of an ointment, and I have met with great success from its use in this manner.

The system I pursue is the following: I have the breast suspended in a sling to prevent all dragging, and pressure exerted on it by means of folded napkins. I then order a free inunction of the iodide of potassium ointment three times a day, administering purgatives internally. For the first two or three days, should it be necessary, I have the milk drawn off once daily by the nurse, and find almost invariably that after a few days all troublesome symptoms pass away, and any anxiety on the score of the milk is removed.