

- Post-partum hæmorrhage, traumatic, 435, 436
- Post-partum prophylactic douching, 6—8
- Postural treatment in oblique presentation, 246—247
- — in prolapse of the cord, 430
- Prague method of delivering after-coming head, 234—235
- Precipitate labour, 366
- Pregnancy, definition of, 80
- diagnosis of, 94—102
- duration of, 80
- extra-uterine, 330—343
- hygiene of, 105—107
- kidney, 263, 264
- multiple, 249—253
- phenomena of, 80—93
- Premature labour, 108, 327
- — induction of, 505—508
- — — in contracted pelvis, 403, 407—409
- Presentation of fœtus, 41—45
- anterior fontanelle, 216—218
- breech, 220—238
- brow, 209—216
- diagnosis of, 50—71
- face, 198—208
- posterior fontanelle, 218, 219
- transverse, 239—248
- vertex, 141—163
- Probable signs of pregnancy, 101, 102
- Prolapse of cord, 428—433
- of limb during labour, 418—422
- Prolapsus uteri during pregnancy, 293
- Prolonged labour, symptoms of, 137—139
- Prophylactic douching, 6—8
- Pruritus vulvæ during pregnancy, 271, 272
- Pseudo-eyes, 102
- Pseudo-osteomalacic pelvis, 392, 393
- Pubiotomy, 540—546
- Puerperal insanity, 493—496
- state, 254—261
- ulcer, 465, 468
- Pulmonary embolus, 484, 485
- Pulse during puerperium, 257
- Purgatives before labour, 173
- during puerperium, 260
- in eclampsia, 309—310
- Pyæmia, 481—483
- Pyrosis of pregnancy, 271
- Quadruplets, 249
- Quickening, 103
- Quintuplets, 249
- Rachitic flat pelvis, 374, 379—381, 401
- Regions of fœtal head, 34
- Relapsing kidney of pregnancy, 264
- Reposition of prolapsed cord, 430, 431
- Respiration, maternal, in pregnancy, 92
- Retained placenta, 369—372
- Retention of urine, in infant, 583, 584
- in pregnancy, 266—268
- Retraction ring, 122—125
- — as a contra-indication to version, 531
- — as a sign of impending rupture, 139, 453, 454
- Retroflexion of gravid uterus, 289—293
- Retro-uterine hæmatocele, 337
- Rheinstädter's flushing curette, 320