

DEPARTMENT OF THE INTERIOR—OFFICE OF THE U. S. ENTOMOLOGICAL COMMISSION.

Providence, R. I., June, 1879.

DEAR SIR,—The Commission desires your co-operation in obtaining facts concerning the habits of the Hessian Fly, with statistics of losses occasioned in your town or county by its attacks; and accounts of the remedies best calculated to prevent its increase, and to destroy it. In brief, the habits of the Hessian Fly are as follows: In May and June two or three small, reddish-white maggots may be found embedded in the crown of the roots of the wheat, at or near the surface of the soil, causing the stalks and leaves to wither and die; the maggots harden, turn brown, then resembling a flax-seed, and change into little *black* midges with *smoky* wings, half the size of a mosquito, which appear in spring and autumn, and lay from twenty to thirty eggs in a crease in the leaf of the young plant. Specimens of the fly may be obtained by sweeping the wheat when three or four inches high, with a gauze net. Please send me specimens of the fly, eggs, maggot and "flax-seed," in vials of alcohol, with notes as to the date when found, and full information as to the insect enemies and parasites.

The *Wheat Midge* is apt to be confounded with the Hessian Fly. It is a small, mosquito-like fly, *orange yellow*, with clear wings, which hovers over fields of young wheat in June. It attacks the heads of the wheat, laying its eggs when the wheat is in blossom. On hatching, the maggots crowd around the young kernels of wheat, causing them to become shrivelled. The maggots in July and August descend into the ground, spinning a round cocoon smaller than a mustard seed, remaining an inch below the surface till the following June.

Information regarding the following topics is respectfully solicited; to be forwarded at the close of the season:

1. When, where, and how are the eggs deposited?
2. When does the maggot appear?
3. When is the "flax-seed" state of the Hessian Fly, or the seed-like cocoons of the Wheat Midge assumed?
4. At what date do the Midges appear in spring and autumn?
5. Look for minute parasites in the eggs and maggots. They may be bred by placing the eggs and maggots with the wheat in bottles covered with gauze, and the parasitic flies preserved in vials of alcohol.