

Males.

- Flagellum denticulate beneath; orbits yellow, except at summit behind *luteoloides*.
 Flagellum ordinary; orbits yellow below I.
 1. Band on segment 1 interrupted; flagellum submoniliform, fulvous beneath *dentariae*, sp. nov.
 Band on segment 1 continuous; flagellum unusually dark, piceous beneath, the joints unusually cylindrical *luteolum*.

Nomada Scop.

Females.

- Head and thorax black, with ferruginous ornaments; abdomen black, with yellow ornaments; interrupted line on segment 1, lateral marks on 2-3, continuous fasciae on 4-5 *vicina*.
 Head and thorax red, sutures, depressed and concealed portions black I.
 1. Scutel quite low, convex, hardly bilobed; joint 4 shorter than 12; black colour of head and thorax rather preponderating over the red; a yellow spot on each side of segments 2-3, two spots on each side of 4, a band or two spots on 5 *simplex*.
 Scutel crested, bilobed 2.
 2. Joint 4 shorter than 12 5.
 Joint 4 as long as 12 3.
 3. Front coxae with short spines; pygidium subacute; yellow fascia on segment 5 opaque, finely rugose, rather sparsely, feebly punctured; scutel strongly crested *denticulata*.
 Front coxae without spines; pygidium broadly subtruncate 4.
 4. A spot on each side of segments 2-3, two subdiscal cuneate spots on 4; band on 5 shining, coarsely punctured; larger *Cressonii*.
 A spot on each side of segments 2-3; smaller *Sayi*.
 5. A spot on each side of segments 2-3, and usually a band or two spots on 5; pygidium broadly rounded, closely pubescent *Illinoensis*.
 A spot on each side of segments 2-5; the smallest species *parva*.

Males.

- Abdomen mainly reddish; vein *a* before *V*₂ 3
 Abdomen mainly black I.
 1. Segments 2-3 with a spot on each side, 1 usually with an interrupted band, 4 with a band or two spots on each side, 5 with a discal band