SYNOPSIS OF BEFS OF OREGON, WASHINGTON, BRITISH COLUMBIA AND VANCOUVER.—III.

BY H. L. VIERECK, ASSISTED BY T. D. A. COCKERELL, E. S. G. TITUS, J. C. CRAWFORD, JR., AND M. H. SWENK.

(Continued from page 196.)

Andrena trachandrenoides, n. sp.

and of 10 mm. Presumably from British Columbia. A very remarkable species, with ochreous pubescence and legs partly pale. If it were not for the character of the antennae, this would have to be placed in

Andrena Kincaidii, Ckll., Proc. Nat. Sci., Phila., 1897, p. 35.

Q, Pullman, Wash. (C. V. Piper); &, and, 10th June, 1895, Olympia, Wash. (T. Kincaid); 7th June, 1895, Seattle, Wash.; 20th May, 1896. Livingston, Vanc.; Q, 6th April, 5th July, 1898; &, 2nd June, 1898; 5th June, 1897; 26th Sept., 1899, Corvallis, Or. (Cordley).

Andrena Kincaidii, Ckll., var. Pascoensis, Ckll., Ent., Lond., 1897, p. 305, Ç, 25th May, 1896, Pasco, Wash. (T. Kincaid), Oregon. Andrena Vernoni, n. sp.

Q 13.5 mm., & 12 mm. Very like Kincaidii, but the dull sculpture separates this at a glance.

Type locality: Vernon, British Columbia. Type Acad. Nat. Sci. Phila.

Andrena Cressoni, Robt., Tr. Am. Ent. Soc., XVIII., p. 56.

Q, 27th May, 1897; 8th June; 4th, 5th, 8th, 11th June, 1898; 4th, 27th April; 12th May, 7th, 10th, 14th June, 1899, Corvallis, Or. (Cordley); o, 11th, 12th, 31st May, 1903, Vernon, B. C.; 29th May, 5th July, Andrena pulverulenta, n. sp.

9 10 mm. The dull fine sculpture, the gray and sericeous pubescence, make of this a sharply defined species.

Type locality: Corvallis, Oregon. Type Acad. Nat. Sci., Phila, 25th Sept., 1898; 3, 15th May, 1897 (Cordley).

Andrena seminigra, n. sp.

2 12 mm. A species with gray pubescence, the abdomen nearly bare and with a steel-blue reflection.

Type locality: Corvallis, Oregon. Type Acad. Nat. Sci., Phila.

Q, 4th May, 1899; 12th, 18th May, 1898; 7th, 9th June, 1898; 3rd, 10th June, 1899, Corvallis, Or. (Cordley).