The Cicadellidæ is represented in America north of Mexico, by 70 genera through which are distributed some 773 species now held to be distinct. On the whole, very little has been done to determine the fauna of any given locality. Several lists, however, have been published and for at least six states the family has been collected extensively and recorded. There may be mentioned the New York and Maine lists by Prof. Osborn, the Tennessee list by De Long, the Wisconsin list by Saunders and De Long, the Missouri list by Gibson and Cogan, and the South Carolina list of Lathrop. In addition to these more extensive lists there have been several minor lists, which need not be mentioned here.

Of the 773 recognized species, some 294 have been recorded from but one locality, the locality of the type and apparently never recovered. The localities of these follow:

of these follow:	The localities
"Am. Sept." 1	T
Boreal Am	Tenn
Carolina	Wisc
Carolina 1	Iowa 4
<b>3.              </b>	Arkansas 1
N. H 1	Tevas
N. Y	Texas
N. J	Kansas
	Nevada 3
•	Utah
D. C	Colorado 62
Va 2	N Movies 62
N. C	N. Mexico
S. C	Arizona 15
	Calif70
DI 3	Washington
20	Oregon 4
Penn 4	Canada 4
Ohio	Canada
III	Manitoba 1
Mich 5	Vanc. Isd
Mich 1	Alaska
Kentucky	Greenland 1
Total-	204
The number of species in addition to	201.

The number of species in addition to the above list that have been recorded from certain of the General Geographical regions may be summed up in the following list:

Generally	distribu	ted throughout the U.S. and C.	
Generally	distribu	ted throughout the U. S. and Canada	32
Generally	through	ted throughout the U. S	24
Generally	through	Atlantic States	33
Occurring	generall	Northern States.	72
"	generan	Jack of the Rocky Mountains	
**	44	smodgi the Southern States.	
"		Wildle Western States or great 1.	
		or the Miss. Myer	
	"		
	"	in the South West	50
"	"	in the North West.	27
"	44	in the Pacific slope.	3
		a wome stope	0.1