

8297. *Geoderces melanothrix*, Kirby.—Under chips on the ground in great numbers and seldom on trees. (See 8288.)
8353. *Trichalophus didymus*, Lec.—Not common. Under logs and in rotten grass.
8470. *Emphyastes fucicola*, Mann.—Occurs but seldom, but then in great numbers. Under logs half buried in sand, during spring.
8471. *Plinthus carinatus*, Boh.—Our commonest weevil. Under logs.
8473. *Pissodes costatus*, Mann.—Not common. Beaten from spruce in April.
8598. *Phycocætes testaceus*, Lec.—Not rare. Under drift-logs on beach in spring.
8599. *Trachodes ptinoides*, Germ.—Common. Under drift-logs on beach in spring. Seen *in cop.*, June 2nd.
8600. *Trachodes 4-tuberculatus*, Mots.—Not common. Found in July. Occasionally in moss at tree roots in winter.
8601. *Trachodes horridus*, Mann.—Common, under logs.
9042. *Elassoptes marinus*, Horn.—Found in extensive colonies under drift-logs in May.
9044. *Rhyncolus brunneus*, Mann.—Not rare; in rotten wood.
Rhyncolus, n. sp.—Rare; under driftwood on sandy beach in May and June.
9074. *Pityophthorus nitidulus*, Mann.—Not rare.
9093. *Xyloterus bivittatus*, Kirby.—Occurs sparingly on newly-cut logs and in moss at tree roots in winter.
9116. *Dryocætes autographus*, Ratz.—Under spruce bark.
9119. *Xylocleptes concinnus*, Mann.—Very numerous in newly-fallen timber.
9142. *Micracis hirtella*, Lec.—Not rare.
9165. *Hylesinus sericeus*, Mann.—Not uncommon in dead bark.
9181. *Dendroctonus rufipennis*, Kirby, var. *obesus* (black form).—Fairly numerous in bark of newly-cut timber.
9187. *Dolurgus pumilus*, Mann.—Common in spruce bark.
Hylurgops glabratus.—Common under spruce.
9373. *Euscaphurus saltator*, Casey.—Common on under side of logs in damp places. [3987. *Dasyllidæ*.]