

- **Tenthredo ferrugineipes*, Cress., Trans. Am. Ent. Soc., Vol. 8, p. 22.
- **Tenthredo flavomarginis*, Nor., Trans. Am. Ent. Soc., Vol. 2, p. 238.
- **Tenthredo mimula*, Cress., Trans. Am. Ent. Soc., Vol. 8, p. 23.
- Tenthredo morosa*, Cress., Trans. Am. Ent. Soc. Vol. 8, p. 23.
- Tenthredo occidentalis*, Cress., Trans. Am. Ent. Soc., Vol. 8, p. 23.
- Tenthredo pallipunctus*, MacGill, CAN. ENT., Vol. 27, p. 282.
- **Tenthredo pectoralis*, Nor., Trans. Am. Ent. Soc., Vol. 2, p. 237.
- **Tenthredo pleuralis*, Cress., Proc. Ent. Soc. Phila., Vol. 4, p. 245.
- **Tenthredo rufipes*, Nor., Proc. Ent. Soc. Phila., Vol. 3, p. 15.
- Tenthredo sectilis*, Cress., Trans. Am. Ent. Soc., Vol. 8, p. 23.
- Tenthredo terminatus*, MacGill, CAN. ENT., Vol. 27, p. 283.
- Tenthredo semirufus*, Nor., Trans. Am. Ent. Soc., Vol. 2, p. 235.
- **Tenthredo variatus*, Nor., Proc. Ent. Soc. Phila., Vol. 3, p. 12.
- Tenthredo variegatus*, Nor., Trans. Am. Ent. Soc., Vol. 2, p. 233.
- **Tenthredo verticalis*, Say, Trans. Am. Ent. Soc., Vol. 2, p. 228.
- Tenthredo ventricus*, MacGill, CAN. ENT., Vol. 27, p. 284.
- **Tenthredo xanthus*, Nor., Proc. Ent. Soc. Phila., Vol. 3, p. 13.
- **Pœcilosoma punctulata*, n. sp.

Descriptions of New Species.

Pœcilosoma punctulata.—Colour black. Length .25 inch. Head narrow from back to front. Eyes set well apart. Abdomen long and extremely flat. In the type specimen it is somewhat concave on the dorsal surface. The base of each segment of the abdomen, except the last two, contains two dorsal white spots, one on each side of the surface, extending nearly half way to the joint in front, and close to the centre of the body. Wings are hyaline, with a beautiful violaceous tinge. Stigma and nervures black.

Emphytus coloradensis.—Colour black. Small species, being only .18 inch in length. Tips of mandibles red, and palpi white. Tibia of front legs, tip of femora, and first joint of tarsus, white, rest of fore legs black. Middle pair of legs have the tibia and femora partly white. Hind legs entirely black. Wings hyaline, nervures black. Lower two-thirds of stigma a lighter colour than upper third.

Blennocampa Gillettei.—Colour black. Length of body .18 inch, to tip of wings .25 inch. Head rounded. Tegulae white. Body short, contracted. All the legs agree in having the tip of femora and tibiae whitish, with the tarsi slightly darker in colour. Wings extend well beyond the tip of the abdomen. Slightly dusky, with beautiful violaceous tinge. Nervures black. Stigma wide, lower two-thirds of lighter colour than upper one-third.