R. Ung. resinos. 3ij.;

Ung. zinci ox. benz., 388.

M. To be applied twice a-day.

Patient may have all kinds of liquid food.

27th.—Cough has entirely disappeared. A more line of granulations, three-fourths of an inch in length. May return home to country. To have ordinary diet after perfect healing of the wound.

July 9th.—Heard indirectly that the child was doing well.

-Boston Med. and Surgical Journal.

OASE OF IMPACTED CALCULUS IN THE URETHRA. EXTERNAL URETHROTOMY. RECOVERY.

BY M. F. GAVIN, M.D., F.R.C.S.L., &C., SURGEON TO OUT-PATIENTS, CITY HOSPITAL, BOSTON.

Henry M., a delicate lad, aged 10 years, born in South Boston and always lived there. Has had the diseases usual to childhood. No pump-water on the premises where he lives. About December, 1868, mother first noticed his water grow cloudy and thick, but free from blood, while the act of urinating caused intense pain in the region of the bladder, extending up towards the lumbar region. From January, 1869, until May of the same year, patient's general health failed, and incontinence of urine and severe "attacks of gravel" became frequent, when he entered the City Hospital under the care of my predecessor. the late Dr. Ropes, who failed to dotect stone. Under treatment and rest he rapidly grew better, and left the hospital, feeling well. For some time after leaving the hospital he remained free from all symptoms of his trouble, except after severe exercise, when the pain in making water returned, and once or twice he suffered from retention, which was relieved by a hot bath.

Early in the autumn the "fits of gravel" returned, grow more severe, lasted longer, and were not relieved by the warm bath. About this time the patient had a very severe homorrhage