

he says, has also been applied here in the form of the spray. He remarks that it is a dangerous drug to put in the hands of an inexperienced person, and, as we have so many other useful remedies for this affection, he thinks it is wise to avoid the use of corrosive sublimate. He has used listerine extensively with good results in the treatment of whooping cough. He employs it in the strength of one drachm to two ounces of water, with an ordinary hand-atomizer, directs the nurse to apply it twelve or more times a day, and finds that little children, even babies, do not object to it. He adds to it tincture of belladonna, potassium carbonate, or ammonia bromide, as the case may demand. Chloride of ammonia he also finds of great service in the form of a spray.—*New York Med. Journal.*

THE TREATMENT OF INTRA-UTERINE DISEASE.—Three papers on this subject (*Brit. Med. Jour.*, Nov. 29, 1855) were read before the Obstetric Section of the British Medical Association, at the fifty-second annual meeting.

Dr. Lombe Athill opened the discussion by a paper in which, after referring to the prejudice which formerly existed against treating the uterine mucous membrane as similar surfaces in other situations are treated, a prejudice which is fast disappearing, the author takes up the consideration of the best method of making such applications as may be necessary. He briefly sums up the conditions which demand such treatment, thus: "all affections of local origin giving rise to profuse menstruation, metrorrhagia, or uterine catarrh, or in which hyperæsthesia of the nerves distributed over the inner surface of the uterus exist." Polypi and other tumors he would exclude, but certain forms of recurrent growths, and of malignant diseases, are to be included. He specifies "local origin" in order to guard against conditions dependent upon affections of the tubes and ovaries being treated in this manner.

The agents he would employ are borax, iodine, carbolic acid, iodized phenol, nitric acid, iodoform, and solid nitrate of silver. Of these remedies, the author finds he uses carbolic acid in nearly seventy per cent. of cases requiring intra-uterine medication, iodized phenol in about fifteen per cent., and nitric acid in three per cent. Borax in the form of a saturated solution in glycerine he uses very occasionally in exceedingly mild forms of catarrh, which have nearly yielded to harsher remedies. Iodine, either in form of tincture or liniment, he considers very inert, as probably very little is absorbed, and its action is mainly caustic. Carbolic acid is the most useful agent, both for curing catarrhs and relieving pain. It should be applied every three or four days, and its use continued some weeks. Iodized phenol, which is iodine dissolved in carbolic acid, one part to three or four,

is of great value in certain cases, especially of hæmorrhage. Nitric acid is the most active of the agents. It does not cause much pain, but the cervical canal should always be protected by a tube of vulcanite or platinum. Iodoform is used in the form of slender crayons in some cases of dysmenorrhœa, and of fetid discharge from the uterus. The solid nitrate of silver as recommended by Sir James Simpson for menorrhagia, depending on subinvolution of the uterus, does not act very satisfactorily, and the author rarely uses it.

All of the agents except iodoform and solid nitrate of silver are used by winding a bit of cotton on a flexible probe, and passing it into the cavity of the uterus. By using dry cotton first, and so removing the mucus, the application will reach the uterine mucous membrane more thoroughly.

There are four conditions where it is necessary to adopt other means: (1) When the mucous membrane is so vascular that the introduction of the probes is followed by hæmorrhage. (2) When the cavity is of large size. (3) When hæmorrhage occurs as the result of vascular growths. (4) When epithelioma affects the cavity of the uterus. For all these conditions the author has been in the habit of dilating, curetting, and applying nitric acid with very good results except in cases of epithelioma. In four cases, the details of which he gives in full, where there was a vascular growth inside the uterus, and where nitric acid failed to control the hæmorrhage, he was induced to try injections of iodized phenol, diluted with an equal part of alcohol, a method which was successful in restraining the hæmorrhage, and prolonging the patients' lives.

Dr. Thomas More Madden, in his paper on the same subject, after a short historical introduction, speaks of the methods of dilating the uterine canal, as by graduated series of dilators, where the tissues are lax, or by tents, especially laminaria, when there is more rigidity. Oftentimes to get the best effect of the agent applied, even when there is no abnormal growth, a moderate curetting beforehand is of advantage. The majority of cases calling for intra-uterine medication, are, according to Dr. Madden, those of what we should call areolar hyperplasia and subinvolution of the uterus. For this treatment he recommends fuming nitric acid applied with the necessary precautions, after thorough dilatation of the canal. Milder applications of carbolic acid, iodoform, and tincture of iodine are recommended later.

For the treatment of tumors within the cavity of the uterus, if submucous or pedunculated, he advises enucleation or écrasement. Within the past few years he has removed forty-two such tumors with thirty-nine recoveries and three deaths. He ends with urging the importance of general as well as local treatment in these cases.

Dr. John W. Byers emphasizes the importance