

Meetings.

MEDICO-CHIRURGICAL SOCIETY OF MONTREAL.

Stated Meeting, April 2nd, 1886.

GEO. WILKINS, M.D., 2ND VICE-PRESIDENT, IN THE CHAIR.

Primary Cancer of Pancreas, with secondary deposits in other organs.—DR. ROWELL exhibited the specimen, and DR. ARMSTRONG related the clinical history of the case :

Mrs. M., aged 80, widow, enjoyed good health until three years ago. Father and two brothers are said to have died of cancer. Admitted to Western Hospital in December, 1885, suffering from loss of appetite, pain after eating, and vomiting. On examination, a hard, round, circumscribed lump, about the size of an orange, was found occupying the epigastrium, just over the region of the pyloric end of the stomach. As little was to be gained from medical treatment, a mixture containing bismuth, hydrocyanic acid and mucilage was prescribed, and she was removed by her friends to her home. It was learned at the time of her death that since her removal from the hospital the vomiting had continued persistently, the most bland liquids, even water, being immediately regurgitated. She had also suffered much pain, for which she had taken morphia pills. Nothing passed bowels for two weeks before death, and she became distinctly jaundiced. At the post-mortem examination, 36 hours after death, only the abdominal cavity was examined. On opening the abdomen, the omentum was found adherent to the anterior abdominal wall. Liver very much enlarged, extending down to level of umbilicus, and containing several large cancerous nodules. Gall-bladder much distended, containing eight ounces of bile and a dark-colored gall stone the size of a cherry. Upon raising the liver, the head of the pancreas was found to be occupied by a cancerous mass, and the surrounding tissues were infiltrated and adherent to it. The walls of the stomach were free from disease. Complete obstruction of the duodenum occurred four inches from the pylorus, caused by pressure of this cancerous mass, together with the adherent and infiltrated tissues about it. A number of the mesenteric glands were also involved. Intestines empty. Spleen slightly enlarged.

New Method for the Relief of Ruptured Perineum.

—DR. TRENHOLME read a paper on this subject, exhibiting drawings of the new method as follows : —This disease must be as old as parturition itself, and yet, beyond the adjustment of the parts, binding the knees together, in recent cases no really successful advance had been made for its cure till the late ever-lamented Dr. Sims introduced his silver suture. The operations of Baker Brown and others were not of any real value, and perhaps the cause or nature of failure was not fully brought out till Emmet's paper upon this subject was given to the world. Now, I do not propose to go over the many points connected with this trouble and the operations attempted for its cure. How much progress has been made can hardly be conceived of by those who have graduated during the last twenty-five years. One of the best and most esteemed surgeons of this city, and, I might say, of this country, endeavored to dissuade a confrere from attempting the operation, stating that "it was sure to be a failure." Not only did he do this, but used his endeavors to prevent the lady from having the operation performed. Thanks, however, to the silver suture and the courage of the operator, the operation was successfully performed and the patient cured. This, occurring in our good city, speaks volumes. For my own part, I think the evils resulting from severe lacerations are very great, and if anything I may say will direct more attention to the prevention of these evils, I will be satisfied. I feel confident that the sum-total of the sorrow and misery arising from this cause vastly exceeds our conception. It is a recognized factor in the causation of subinvolution of the vagina and uterus, and I am persuaded its results are not limited to these organs, but that the tubes and round ligaments share in the same mischief. It is a fruitful cause of retro-luxations of the uterus and prolapsus of bladder. Of all the marital misery and personal distress I need say nothing; these, of course, vary with the peculiarities of individual cases and the extent of the disease. I will not speak of the well-known preparation of the patient required, especially in extensive lacerations; you all know as to this and the after-treatment also. There is one remark I wish to make as to what is known as the perineal body. Some writers have made light of its existence, because its anatomy and relations are not sufficiently definite to merit, as they think, this appellation. That every uninjured perineum has such a body is unquestionable,