WORMS (Continued) .--

B. Ol. filicis maris ..... f3iij.
Ol. chenopodii...... f3 j.
Ol. terebinth ...... f3 ij.
Emul. ol. ricini (50 per cent.) q. s. ad ..... f3 ij.

M. Sig.: Teaspoonful twice a day for a child of six years. (Tapeworm). -L. Starr,

B. Hydrarg. chlor. mit... gr. j. Resinæ jalapæ..... gr. ij. Pulv. scammonii.... gr. v.

M. Et ft. chart. No. i. Sig.: To be taken at bedtime for a child of six years. (Seatworm.)—Goodhart and Starr.

B. Tanret's pelletierini... I bottle.

Sig.: In the evening use a large laxative injection and take only milk.

The next morning mix the contents of a bottle with a glass of water, and take at one dose; one hour after, take one ounce of compound tincture of jalap mixed with a half glass of water. (Tapeworm.)—L. Starr.

After a light diet the evening before, give the following on an empty stomach :—

M. Sig.: Take in two doses, half an hour apart. (Tapeworm.)—*Phar*mas Zeit.

 I? Tr. ferri chlor..... f 3 ss. Aquæ...... Oj.

Sig.: Inject one-fourth to one-third. (Seatworms.)—*Ringer.* 



IS PRONOUNCED BY ALL PHYSICIANS who have examined it, and patients who have used it to be the best and most perfect fitting supporter made. It is self-adjusting and affords instant relief.

