

had quite normal pressure (110) which was not definitely influenced by suprarenal powder.

*Acute Articular Rheumatism involving the heart, 8 cases:*—Six normal, 2 slightly subnormal.

*Acute Articular Rheumatism, heart not affected, 5 cases:*—One normal, 4 slightly subnormal.

*Chronic Articular Rheumatism, 4 cases:*—All normal.

*Gonorrhœal Rheumatism, 8 cases:*—Six normal, two moderately subnormal.

*Rheumatoid Arthritis, 10 cases:*—Six normal, and one of 85, one 95, one 100, one 105.

*Gout, 4 cases:*—Only one of them well marked. This one had albuminuria and moderate arterio-sclerosis, yet a reading scarcely over the normal (130). The others had normal pressure.

*Hemiplegia, 1 case:*—In the course of chronic nephritis, with heart negative and a moderate degree of arterio-sclerosis, showed high pressure in both hands, but 10 mm. lower in the paralyzed than in the sound hand. Another similar case with higher pressure gave equal readings in both hands. Two other hemiplegias with slightly low pressure had practically equal readings in both hands.

*Tabes Dorsalis, 14 cases:*—Eleven normal, 3 slightly raised, one of the latter having albuminuria, and two arterio-sclerosis.

*Cerebral Tumour, 8 cases:*—All normal pressure; one who gave a slightly high reading, but was suffering intense pain at the time of observation.

*General Paralysis of the Insane, 1 case:*—With moderately high tension, slight arterio-sclerosis, urine 1012, trace of albumin, normal.

*Friedreich's Ataxia, 2 cases:*—One with albuminuria and small amount of sugar in urine, slightly raised (140); one combined with cerebellar ataxia, normal.

*Acute Ascending Paralysis, 1 case:*—Involving phrenics, in a male, aged 27. Pressure was slightly raised (140) during worst phases of attack, and remained so during several weeks, but gradually came down and was nearly normal at time of discharge; no albuminuria.