

SAXIFRAGA TOLMÆI, T. & G.

The range of this species in Canada is extended by the following references: Mountain summits opposite Vancouver, B.C. (*T. Henry.*) North of Ch'liiwack River, south slope of Cheam Range, B.C. alt. 7,000 feet. No. 34,948; Tami Hy Mountain, Chilliwick Valley, B.C. No. 34,949. (*J. M. Macoun.*)

HEMIEVA VIOLACEA, Wheelock.

Near Kaslo, B.C., 1901. (*Miss Ethel Blackman.*) New to Canada.

MITELLA TRIFIDA, Graham.

Alpine slopes, Mount Arrowsmith, Vancouver Island, 1887. (*John Macoun.*) Same locality, 1901. (*J. R. Anderson* and *Dr. James Fletcher.*) Common between 4,500 and 5,500 feet altitude in the Chilliwick Valley, B.C. Nos. 34,864 and 34,865. (*J. M. Macoun.*)

MITELLA BREWERI, Gray.

Mountains at Chilliwick Lake, B.C. No. 34,866. (*J. M. Macoun.*) Western limit.

MITELLA OVALIS, Greene.

Damp places, Goldstream, Vancouver Island, 1887. No. 8,497. (*John Macoun.*) Our only Canadian specimen.

MITELLA CAULESCENS, Nutt.

A common plant in woods throughout the Chilliwick Valley, B.C. Nos. 34,860, 34,861 and 34,863. (*J. M. Macoun.*) Our first Canadian specimens.

MITELLA PENTANDRA, Hook.

Alpine woods, Mt. Arrowsmith, Vancouver Island, 1887. (*John Macoun.*) Same locality, 1901. (*J. R. Anderson* and *Dr. James Fletcher.*)

ONAGRA OAKESIANA, Britton.

Gaspé, Gulf of St. Lawrence; near London, Ont.; Port Colborne, Ont., No. 44,465; Southampton, Ont., No. 44,464. (*John Macoun.*) Seldom separated from *O. biennis*.