Co., Ont., in the summer of 1897, and again in 1898. Communicated by Dr. T. J. W. Burgess.

LESPEDIZA VIRGINICA (L.) Britt.

In thickets at Leamington, Ont., 1892. (John Macoun.) New to Canada.

LESPEDEZA FRUTESCENS, Britt

L. Stuvei, var. intermedius, Gray. Man. Ed. vi. p. 147.

L. reticulata, Macoun, Cat. Can. Plants, vol. 1, pp. 119 and 511.

Not rare in sandy woods and thickets from Niagara to Sarnia.

LATHYRUS MYRTIFOLIUS, Muhl

L. paluster var, myrt folius, Gr.; Macoun, Cat. Can. Plants, vol. I, p. 122.

In thickets at Brandon. Man. Herb. No. 12,528. 1896. (John Mocoun.) Western limit.

PRUNUS AVIUM, L,

Along a creek bank west of Niagara-on-the-Lake, Ont. (J. Pewness) Not before recorded in Canada.

SPIRÆA LOBATA, Jacq.

Escaped from cultivation and naturalized at Boylston, N.S. (Dr. C. A. Hamilton.) Not before recorded in Canada.

SPIRÆA SORBIFOLIA, L.

Along roadsides near Biddeck, Cape Breton Island, N.S., 1898. (John Macoun.) Escaped from cultivation, but not before recorded as well naturalized.

POTENTILLY NEMORYLIS, Nestler: Macous, Cat. Can. Plants, vol. 1, p. 144.

Discovered many years ago at Ste. Ann's, Cape Breton Island, N.S., by Dr. Lawson. Collected in 1898 by Prof. Macoun between Ste. Ann's and Baddeck Bay, Cape Breton Island.

RIBES FLORIDUM, L'Her.; Micoun, Cat. Cin. Pioits, vol. 1, 0, 153

Collections of recent years have shown this process to be more widely distributed west of Manitoba than has supposed.