

STRAWBERRY
 Leaf-spot and white grub.
 For leaf spot set out only healthy plants with no sign of disease. First season spray with D before blossoms open and keep plants covered with mixture throughout the season. Second year spray before blossoming with D, and again soon after picking; or mow and burn over after picking. Don't take more than two crops off. Plow down at once after second crop. For white grubs dig out as soon as injury is noticed. Do not plant on land broken up from old meadow or pasture for three years after breaking.

BEAN
 Anthracnose and bacterial blight.
 Get seeds from pods showing no signs of disease. Do not work among the plants if they are wet with rain or dew. Spraying scarcely pays as a rule.

CABBAGE & TURNIP
 Flea-beetles, caterpillars, root maggots, aphids.
 For flea-beetle on turnip sow after June 21st, or dust plants as soon as they appear above ground with Paris Green, or spray with Bordeaux and a poison. Repeat in two or three days. For caterpillars dust with Paris Green until heads begin to form on cabbage and cauliflower, then spray with catnip-pyrethrum, 1 ounce to 2 gallons water. For root maggots use medium thick tarred felt paper discs, putting on as soon as plants are set out, or set out plants after July 1st. For aphids use kerosene emulsion as soon as they appear.

POTATO
 Tip burn, early blight, late blight, scab, Colorado beetle, flea-beetle, tubers.
 Keep foliage covered with D from time plants are about 5 inches high. Take special precautions to see this is well done if weather is at all damp after about 15th July, as late blight begins about this time. Add a poison to each application when necessary. For scab, soak tubers before cutting 2 hours in formalin solution, 1 pint of formalin to 30 gallons of water. Spread out on grass to dry. Wash all boxes, bags or other vessels to be used in same liquid. Plant none but perfectly healthy tubers.

TOMATO
 Leaf blight, black rot, fruit, flea-beetle.
 Spray plants in seed bed with D. Keep foliage in field covered with D until danger of staining flea-beetle.

ASPARAGUS.
 Rust, beetles.
 For rust, let no plants, not even wild ones, mature during cutting season. Late in fall, when growth is about over, cut and burn old plants. For beetles let poultry run in the plantation. After cutting season is over spray with arsenate of lead; repeat in two weeks. May add sticker and a little lime.

NOTE.—A1—Concentrated lime-sulphur, strength 1.030 specific gravity (1.032 for San José scale).
 A2—Concentrated lime-sulphur, strength 1.099 specific gravity.
 A3—Concentrated lime-sulphur, strength 1.008 specific gravity.
 B—The old home-boiled lime-sulphur, 20.15.40 formula.
 C—Self-boiled lime-sulphur.
 D—Bordeaux mixture, 4.4.40 formula.