Eurymus nastes (Boisd.) = Colias nastes of former list. I have not reexamined specimens, as they are apparently no longer extant.

Pamphila comma (Linn) = Hesperia comma of my former list. The single specimen obtained was not examined by me in my study of the species of Pamphila (Memoirs Bost. Soc. Nat. Hist. ii., 341), and is the only specimen I have seen of P. comma from America. It belongs to the var. catena Staud, found in northern Scandinavia and Lapland, and exactly resembles the specimen of that variety figured by me in the memoir referred to above. Moschler has already noted that it is this variety which occurs in Labrador.

Hesperia centaureae Ramb. Nothing to be added.

HYMENOPTERA.

Urocerus flavicornis Fabr. Common on Caribou Island. " cyaneus Fabr. Hopedale.

Euura orbitalis Norton. Var. a. b. Caribou Island. Nematus Labradoris Norton. Caribou Island.

"	malacus	"	**	"
•6	fallax	**	••	""
"	monela	.6	••	
46	fulvipes	••	"	**
**	placentus	"	••	٠٠
Allantus originalis		**	••	**

Macrophya (Pachyprotasis) omega Norton. Caribou Island. Tenthredo mellinus Norton. Caribou Island.

" cinctitibiis " " "

Formica herculanea Linn. Whole coast.

" sanguinea Latr. Straits of Belle Isle.

Vespa maculata Linn. Southern coast, Mecatina Island. norvegica Fabr. Caribou Island.

Bimbus lacustris Cresson. Whole northern coast; common.

" kirbyellus Curtis. Sloop Harbor and Hopedale.

" frigidus Smith. Square Island and Hopedale.

" nivalis Dahlb. Caribou Island and whole coast northward.