

Type Am. Ent. Soc., Phil.

Type locality, Vancouver I., Canada.

Species of *Triepeolus*, related to *occidentalis*, and represented only by the ♂:

First joint of the flagellum not much more than one-half the length of the second 1.

1. Legs and labrum ferruginous *occidentalis*, var. 1.

First joint of the flagellum nearly as long as the second 2.

2. Femora, except anterior pair, largely black; anterior femora, all tibiae and all tarsi ferruginous; labrum reddish *occidentalis*, var. 2?

Legs entirely black, excepting the pubescence and the tarsi, the former being white and the latter brownish n. sp.?

Triepeolus occidentalis (Cress.).—Tr. A. E. S., VII, p. 87, 1879, ♀ ♂.

The co-types of this species are from Colorado, and have an almost entirely black labrum.

Var. 1. Oregon. (Received through the courtesy of Mr. J. H. Lovell.)

Triepeolus occidentalis, var. 2?

Oregon. (Through Mr. J. H. Lovell.)

Triepeolus, n. sp.?

Vernon, B. C., 12th Aug., 1904 (Harvey).

Epeolus, Latr.

Markings of abdomen white or pale or cream colour; mark on first abdominal segment a transverse band; bands on second to fourth segments interrupted in the middle line; size small; femora

black *olympiellus*.

Epeolus olympiellus, Ckll.—Ann. & Mag., N. H., XIII, p. 41, 1904.

Olympia, Wash., 2nd July, 1896 (Kincaid).

Epeolus tristicolor, Vier., n. sp.

Related to *autumnalis*, from which it differs in size, structure and colour.

The italicized characters in *T. penepectoralis* occur in this species, and are not here repeated.

♀.—8.5 mm. Frontal keel originating as in *T. penepectoralis*, but not extending higher than the middle of the front even as a raised line, and not connected with the anterior ocellus by a shining line; clypeus and supraclypeal space sculptured nearly like the cheeks, somewhat shining; labrum sculptured much like the front, with two short teeth on